



# Transmission of potato spindle tuber viroid between *Phytophthora infestans* and host plants

O.S. Afanasenko , A.V. Khiutti, N.V. Mironenko, N.M. Lashina


All-Russian Research Institute of Plant Protection, Pushkin, St. Petersburg, Russia  
 olga.s.afan@gmail.com


**Abstract.** Potato spindle tuber viroid (PSTVd) is a naked, circular, single-stranded RNA (356–363 nucleotides in length) which lacks any protein-coding sequences. It is an economically important pathogen and is classified as a high-risk plant quarantine disease. Moreover, it is known that PSTVd is mechanically transmitted by vegetative plant propagation through infected pollen, and by aphids. The aim of this study is to determine the possibility of viroid transmission by potato pathogen *Phytophthora infestans* (Mont.) de Bary. PSTVd-infected (strain VP87) potato cultivars Gala, Colomba, and Riviera were inoculated with *P. infestans* isolate PiVZR18, and in 7 days, after the appearance of symptoms, re-isolation of *P. infestans* on rye agar was conducted. RT-PCR diagnostics of PSTVd in a mixture of mycelia and sporangia were positive after 14 days of cultivation on rye agar. The PSTVd-infected *P. infestans* isolate PiVZR18v+ was used to inoculate the healthy, viroid-free plants of potato cv. Gala and tomato cv. Zagadka. After 60 days, an amplification fragment of PSTVd was detected in the tissues of one plant of tomato cv. Zagadka by RT-PCR with the primer set P3/P4, indicating successful transmission of PSTVd by *P. infestans* isolate PiVZR18v+. This result was confirmed by sequencing of the RT-PCR amplicon with primers P3/P4. The partial sequence of this amplicon was identical (99.5 %) to PSTVd strain VP87. RT-PCR showed the possibility of viroid stability in a pure culture of *P. infestans* isolate PiVZR18v+ after three consecutive passages on rye agar. PSTVd was not detected after the eighth passage on rye agar in *P. infestans* subculture. These results are initial evidence of potato viroid PSTVd being bidirectionally transferred between *P. infestans* and host plants.

Key words: potato; tomato; PSTVd strains; transmission; *Phytophthora infestans*; RT-PCR detection.

**For citation:** Afanasenko O.S., Khiutti A.V., Mironenko N.V., Lashina N.M. Transmission of potato spindle tuber viroid between *Phytophthora infestans* and host plants. *Vavilovskii Zhurnal Genetiki i Seleksii = Vavilov Journal of Genetics and Breeding*. 2022;26(3):272-280. DOI 10.18699/VJGB-22-34

# Трансмиссия вириода веретеновидности клубней картофеля между *Phytophthora infestans* и растениями-хозяевами

О.С. Афанасенко , А.В. Хютти, Н.В. Мироненко, Н.М. Лашина

Всероссийский научно-исследовательский институт защиты растений, Пушкин, Санкт-Петербург, Россия  
 olga.s.afan@gmail.com

**Аннотация.** Вириод веретеновидных клубней картофеля (ВВКК) представляет собой кольцевую одноцепочечную РНК длиной 356–363 нуклеотида, в которой отсутствуют какие-либо последовательности, кодирующие белок. ВВКК является экономически значимым заболеванием картофеля, имеющим статус карантинного. Известно, что ВВКК передается механически при вегетативном размножении растений, через инфицированную пыльцу и с помощью тлей. Целью данного исследования было определение возможности передачи ВВКК (штамм VP87) от растений картофеля и томата, зараженных вириодом, патогену *Phytophthora infestans* (Mont.) de Bary и от *P. infestans* – растениям картофеля и томата. Сорта картофеля Гала, Коломба и Ривьера, инфицированные ВВКК, инокулировали изолятом *P. infestans* PiVZR18; через 7 дней после появления симптомов фитофтороза провели повторное выделение *P. infestans* в чистую культуру на ржаной агар. Через 14 дней культивирования *P. infestans* на ржаном агаре в смеси мицелия и спорангиев методом ОТ-ПЦР был обнаружен ВВКК. ВВКК-инфицированным изолятом *P. infestans* (PiPSTVdv+) провели инокуляцию растений томата сорта Загадка и растений картофеля сорта Гала. Через 60 дней в листьях томата сорта Загадка методом ОТ-ПЦР с праймерами P3/P4 был выявлен диагностический продукт амплификации 360 п.о., свидетельствующий об успешной трансмиссии ВВКК изолятом PiVZR18v+. Результаты были подтверждены секвенированием продукта амплификации ВВКК. Нуклеотидная последовательность вириода в растении томата, зараженном изолятом PiVZR18v+, оказалась на 99.5 % идентичной использованному в эксперименте штамму VP87. Для доказательства возможности сохранения вириода в чистой культуре *P. infestans* изолят PiVZR18v+ пассировали на ржаном агаре с интервалом 30 дней. После трех последовательных пассажей на ржаном агаре ВВКК был диагностирован в культуре изолята, что подтверждено секвенированием продукта

амплификации с вириод-специфичными праймерами. В субкультуре *P. infestans* после восьмого пассажа на ржаном агаре ВВКК не обнаруживался. Полученные данные свидетельствуют о двунаправленной передаче ВВКК в патосистеме *P. infestans* – растение-хозяин.

Ключевые слова: картофель; томаты; штаммы ВВКК; трансмиссия; *Phytophthora infestans*; ОТ-ПЦР-диагностика.

## Introduction

Potato spindle tuber viroid (PSTVd) is an economically important pathogen, classified as a high-risk plant quarantine disease. According to the European Plant Protection Organization (EPPO), the disease has been reported in 37 countries on all continents (<https://gd.eppo.int/taxon/PSTVD0/distribution>). In Russia, and other former Soviet Union regions, PSTVd was detected in about 50–70 % of *in vitro* potato plants (Kastalyeva et al., 1992).

Natural infections of PSTVd have been found in the field, mainly in solanaceous crops, such as pepino (Puchta et al., 1990), potato (Diener, Raymer, 1969), and tomato (Puchta et al., 1990). Potato plants infected with PSTVd become smaller and show leaf yellowing, and infected tubers become smaller and cracked. The reduction in tuber weight depends on the viroid strain, potato cultivar, and length of time they have been infected with PSTVd (Pfannenstiel, Slack, 1980). Furthermore, a reduction in tuber yield of up to 24 % has been reported in cultivar Saco infected with mild strains of PSTVd, however, severe strain reduced the yield by up to 64 % (Singh R.P., 1970). In addition to direct losses, it is important to take into account indirect losses that can be significant due to the quarantine status.

PSTVd belongs to Family Pospiviroidae (IPOSPP), Genus *Pospiviroid* (IPOSPG) and consists of a naked, circular, single-stranded RNA (356–363 nucleotides in length) – the smallest among plant pathogens lacking a protein-coding ability – therefore, it is a parasite of the host transcription mechanism (Yanagisawa et al., 2019).

PSTVd has a wide host range of at least 138 species across 10 families (Singh R.P., 1973). The main hosts are from the Solanaceae family (Owens et al., 1992; Mertelik et al., 2010; Mackie et al., 2016). PSTVd is transmitted mechanically (Verhoeven, Roenhorst, 2010), by aphids (Syller et al., 1997). Moreover, it was found to be vertically transmitted through pollen to progeny seeds on potato (*Solanum tuberosum*) and tomato (*S. lycopersicum*) and horizontally transmitted through infected pollen to other potato and tomato plants (Kryczyński et al., 1988; Singh R.P. et al., 1992; Matsushita, Yanagisawa, 2018).

Viruses are common in fungi and oomycetes and some of these viruses share sequence identities with plant viruses belonging to different families and genera (Mascia et al., 2019). There are several examples of plant–virus transmission by phytopathogenic fungi and oomycetes. It was shown that soil-inhabiting fungi *Olpidium brassicae* and *O. radiale* belonging to Chytridiales and *Polymyxa graminis*, *Spongospora subterranean*, and *Synchytrium endobioticum* – belonging to the order Plasmodiophorales – transmit plant viruses (Bhat, Rao, 2020). Replications of the tobacco mosaic virus were demonstrated in the phytopathogenic fungi *Colletotrichum acutatum*, *C. clavatum*, and *C. theobromicola* (Mascia et al., 2019), cucumber mosaic virus was reported in *Rhizoctonia solani* (Andika et al., 2017), artichoke Italian latent virus,

artichoke mottled crinkle virus, potato virus X, potato virus Y, tobacco mosaic virus and cucumber mosaic virus plus its satellite RNA can replicate and persist in *Phytophthora infestans* at least through the first subculture (Mascia et al., 2019).

Wei et al. (2019) obtained preliminary data on the possibilities of replicating hop stunt viroid (HSVd), iresine 1 viroid belonging to the Pospiviroidae and avocado sunblotch viroid (Avsunviroidae) in at least one of phytopathogenic ascomycete fungi *Cryphonectria parasitica*, *Valsa mali*, and *Fusarium graminearum*.

Oomycete *P. infestans* causes significant losses to potato and tomato crops on a global scale. Despite intensive use of fungicides, the pathogen is constantly and ubiquitously present in potato crops. *P. infestans* is also the most harmful and widespread tomato pathogen, both in field and greenhouse conditions. This oomycete has a high adaptive potential to the host plants, which may indicate the formation of competitive relationships with other potato and tomato pathogens. In this regard, it is of interest to identify a possible role of *P. infestans* in transmission of PSTVd to potato and tomato plants.

## Materials and methods

**Plant materials.** Potato cultivars that, according to our data, were susceptible to both PSTVd strain VP87 and *P. infestans* – Gala, Riviera, and Colomba and tomato cultivars Zagadka, Moskvich, and Damskiy Palchik were included in the study.

These potato and tomato cultivars were registered in the Russian State Register of Breeding Achievements.

**PSTVd strains.** Two intermediate PSTVd strains, VP35 (GenBank accession no. LC523658) and VP87 (LC523667), and severe strain FP10-13 (LC523676) deposited in the international information database DDBJ (DNA Data Bank of Japan), Data set “Viruses” (<http://blast.ddbj.nig.ac.jp/>) were used in the study. These strains were isolated from infected potato leaves from the Volga Federal District (VP87 and VP35) and tubers from the Far Eastern Federal District (FP10-13) in 2019 (Matsushita et al., 2021).

The PSTVd strains were supported on living tomato plants of Russian cultivars Zagadka, Moskvich, and Damskiy Palchik.

**Isolate of *P. infestans*.** Isolate PiVZR18 of *P. infestans* was used in the experiments on viroid transmission. PiVZR18 was isolated from the natural population of *P. infestans* in the Leningrad Region (northwest of the European part of Russia) in 2018. Eight virulence genes (1, 2, 3, 4, 6, 7, 10, and 11) were identified in this isolate on a set of Black’s differentials (Black et al., 1953).

**Viroid inoculation of plants.** Potato and tomato plants were grown in a growth room at a temperature of 25 °C with a photoperiod of 16h/8h (day/night) in 2l pots filled with “Terra vita” soil. Seven-day germination potato plants and 14-day tomato plants were used for inoculation by PSTVd.

To prepare the inoculum, 0.1 g of fresh tomato leaf tissue – 60 days post inoculation (dpi) with PSTVd strain VP87 – was

ground in 1 ml sodium phosphate buffer (pH 7.0) and filtered through cheesecloth.

For mechanical inoculation, the cotyledons of tomato were dusted with carborundum and gently rubbed over the surface of the leaves with a plastic pestle. Ten microliters of inoculum was placed on the injured leaf surface and rubbed several times with a sterile plastic pestle. The inoculated plants were incubated for two months at 25 °C with light intensity (fluorescent, 40 W, ×4).

At 60 dpi, the presence of viroid in the inoculated tomato plants was determined by RT-PCR.

To inoculate 7-day potato plants of the cultivars Gala, Riviera, and Colomba, a 0.5–1.0 cm longitudinal stem incision was performed with a sterile razor on a stem apex (Suppl. Fig. 1)<sup>1</sup>, and 10 µl of the PSTVd VP87 strain suspension – obtained as described above – was applied. Three plants of each potato cultivar were inoculated and the assay was repeated three times. In 60 dpi, the presence of PSTVd in the inoculated plants of potato cultivars was determined by RT-PCR.

***P. infestans* inoculation of plants.** Isolate PiVZR18 of *P. infestans* was cultured on rye agar medium (1.0 Li ddH<sub>2</sub>O, 60.0 g rye organic berries (grind in blender), 20.0 g sucrose, 15.0 g agar) for 30 days in the dark at 15 °C for propagation and morphological observation (Medina, Platt, 1999).

Before inoculating the plants, the suspension was incubated at 12 °C for 2.5–3 h to release zoospores. Upon RT-PCR detection for viroid infection, both healthy and viroid-infected tomato and potato plants were inoculated with a suspension of *P. infestans* at a concentration of 50.000 zoospores in 1 ml. After inoculation, the plants were placed in humid chambers with a 14 h light period, at 23 °C during the day and 15 °C at night for a period of 13 dpi. To study PSTVd transmission from *P. infestans* to host plants, when typical symptoms of late blight appeared, the humid chamber was removed and the development of *P. infestans* slowed down. The affected leaves were removed and the plants continued to grow. For the purposes of PSTVd diagnostics, the upper leaves of the plants without late blight symptoms were cut off.

**Viroid inoculation of *P. infestans*.** Inoculum of PSTVd was obtained from the infected tomato plants as described above and applied to the 14-day *P. infestans* culture by transferring 10 µl per Petri dish (in the dish center). After inoculation, the culture was left to grow for 15 days at 10 °C. Then, mycelia from the periphery and from the center of the colony were transferred separately to a fresh medium. The culture was left in the same conditions for 30 days, after which RT-PCR analysis was conducted.

**Isolation of *P. infestans* from infected potato and tomato plants.** Seven days post inoculation (dpi) after the symptoms of late blight appeared, *P. infestans* was isolated from the plants. Sections of the infected leaves were placed between tuber slices of the healthy cv. Colomba, and at 6 days, mycelium was transferred with a needle to the surface of rye agar. The isolates were cultured for 30 days at 15 °C in the dark and then transferred to a fresh medium.

**Viroid detection and sequencing.** We collected the uppermost leaves from the inoculated potato or tomato plants

at 60 dpi. PSTVd detection in pure culture of *P. infestans* was carried out after 30 days of growing on rye agar. Approximately 0.1 g of tissue from leaves or mycelium was used for RNA extraction. Total RNA was extracted using the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany) as per the manufacturer's instructions ([http://www.genome.duke.edu/cores/microarray/services/rna-qc/documents/RNeasy\\_Mini\\_Handbook.pdf](http://www.genome.duke.edu/cores/microarray/services/rna-qc/documents/RNeasy_Mini_Handbook.pdf)) and subsequently used for one-step RT-PCR. Primer sets P3/P4 (Behjatnia et al., 1996) and P1/P2 (Gross et al., 1978) or 68PV-R+87PV-F (Yanagisawa et al., 2019) were used to detect PSTVd. RT-PCR was prepared with the PrimerScript One-Step RT-PCR Kit ver 2 reagents in 10 ml (Takara Bio Inc., Shiga, Japan) following the manufacturer's instructions. The primer set ITS4/ITS5 (White et al., 1990; Ristaino et al., 1998) was used to detect the ITS region of *P. infestans* as an internal control.

RT-PCR was carried out on a MyCycler Thermal Cycler (Bio-Rad, California, USA) at 50 °C for 30 min, 94 °C for 2 min, followed by 35 cycles of 94 °C for 30 sec, 60 °C for 30 sec and 72 °C for 30 sec. An additional elongation step was performed at 72 °C for 5 min followed by storage at 12 °C. The sizes of the diagnostic fragments of PSTVd and the ITS region were 360 and 946 bp, respectively.

The PSTVd amplicons were sequenced at the Beagle Company (St. Petersburg, Russia). Alignment and manual editing of nucleotide sequences were performed using Vector NTI Advance 10 software (Thermo Fisher Scientific). The obtained nucleotide sequences were tested for similarity with PSTVd strain VP87 (LC523667), used in this study, deposited in the international information database DDBJ (DNA Data Bank of Japan), data set "Viruses" (<http://blast.ddbj.nig.ac.jp/>).

## Results


There is an absence of data on the possibility of replicating PSTVd in *P. infestans* and bidirectionally transferring it between host plants and *P. infestans*. In this study, we investigated the possibility of PSTVd transmission (1) from host plants to *P. infestans*, (2) from *P. infestans* to host plants, and (3) the possibility of PSTVd stability in pure cultures of *P. infestans*.

### Transmission of PSTVd from host plants to *P. infestans*

**From potato plants.** Upon confirming PSTVd infection in tomato plants of cv. Zagadka by RT-PCR, inoculation of three potato cultivars (Gala, Colomba, and Riviera), using as an inoculum source extracted-sap of tomato infected with PSTVd strain VP87, was conducted (see Suppl. Fig. 1). After 60 days, detection of PSTVd presence in plants of these cultivars was carried out by RT-PCR with the P3/P4 primer set (Fig. 1). The brightest amplicons indicating a high accumulation of the viroid were found for cv. Gala and cv. Colomba. The viroid accumulation was lower in three plants of cv. Riviera (see Fig. 1).

PSTVd-infected and uninfected (control) potato plants were inoculated with *P. infestans* isolate PiVZR18. Seven days after the appearance of symptoms (Fig. 2, a) caused by *P. infestans*, the pathogen was first re-isolated from these cultivars on tuber slices of healthy cv. Colomba and then transferred to pure culture on rye agar (see Fig. 2).

<sup>1</sup> Supplementary Figures 1–3 are available in the online version of the paper: <http://www.bionet.nsc.ru/vogis/download/pict-2022-26/appx5.pdf>



**Fig. 1.** RT-PCR detection of PSTVd with the primer set P3/P4 in leaves of infected potato plants by PSTVd strain VP87 before inoculation with *P. infestans*.

1–3, Individual plants of cv. Riviera; 4, mock-inoculated cv. Riviera; 5–8, individual plants of cv. Colomba; 9, mock-inoculated cv. Colomba; 10–13, individual plants of cv. Gala; 14, mock-inoculated cv. Gala; 15, negative control (distilled water); M, molecular weight marker 50 bp DNA ladder (Primetech DNA Ladder).

After culturing of *P. infestans* isolates on rye agar, PSTVd detection by RT-PCR was conducted with the primer set P3/P4 (Fig. 3, a). Amplicons of ~360 bp – indicating the presence of PSTVd – were detected in a mixture of mycelium and sporangia of pure culture of *P. infestans* isolates after colonization on the cultivars Gala, Riviera, and Colomba (see Fig. 3, a). To control for the negative results of viroid detection that are not due to the quality of RNA extracted from *P. infestans* samples, we used PCR with universal primers ITS4/ITS5 on rDNAs, which are species specific for *P. infestans* and displayed an amplicon of 946 bp (Ristaino et al., 1998) (see Fig. 3, b).


**From tomato plants.** The 15 plants of the two tomato cultivars (Moskvich and Damskiy Palchik) infected with three PSTVd strains (VP87, FP10-13, and VP35) were inoculated with the *P. infestans* isolate PiVZR18 to confirm the results obtained. Seven days after the appearance of the symptoms caused by *P. infestans*, 15 cultures of the pathogen were re-isolated from these plants, first on tuber slices of healthy cv. Colomba. After mycelial overgrowth on the surface of tuber slices, the first detection of viroid presence in the mycelium was performed. Out of 15 isolates, the brightest fragment indicating viroid infection of mycelium was detected with the primer set P3/P4 in the *P. infestans* isolate from cv. Moskvich infected with PSTVd strain VP87 (Fig. 4, line 1). Positive PSTVd detection was also obtained for cv. Damskiy Palchik, infected by PSTVd strain VP87 (line 4) and for the isolates from cv. Moskvich infected by PSTVd strain FP10-13 (lines 6–8) and by strain VP35 (lines 10–13). Weak amplicons were obtained in lines 3, 5, 9, 14, which indicates a low initial concentration of PSTVd in *P. infestans* mycelium obtained from the host plant (see Fig. 4).

#### PSTVd transmission from *P. infestans* to host plants

The PSTVd-infected *P. infestans* isolate PiVZR18v+ was used to inoculate the healthy, viroid-free plants of potato cv. Gala and tomato cv. Zagadka.


After 60 days, an amplification fragment of PSTVd was detected in the tissues of one plant of tomato cv. Zagadka by RT-PCR with the primer set P3/P4, indicating successful transmission of PSTVd by *P. infestans* isolate PiVZR18v+ (Fig. 5).

The detection of PSTVd in the RNA of cv. Zagadka inoculated with the *P. infestans* isolate PiVZR18v+ was



**Fig. 2.** PSTVd transmission from potato to *P. infestans*.

a, Mock-inoculated (distilled water) potato cv. Colomba; b, PSTVd-infected (strain VP87) potato cv. Colomba; c, symptoms of *P. infestans* on PSTVd-infected potato cv. Colomba (strain VP87); d, tuber slices of healthy cv. Colomba inoculated with *P. infestans* from the infected plants; e, *P. infestans* isolates from the PSTVd-infected plants of cv. Colomba on rye agar.




**Fig. 3.** RT-PCR detection of PSTVd in cultures of *P. infestans* isolated from potato plants infected with the PSTVd strain VP87 (a) and control of *P. infestans* RNA presence (b).

a, RT-PCR detection with the primer set P3/P4; b, RT-PCR amplification of the same RNA samples with the primer set ITS4/ITS5. Lines: 1, PSTVd-uninfected *P. infestans* isolate PiVZR18; 2, *P. infestans* isolates from PSTVd-infected potato cv. Gala; 3, 4, from cv. Riviera; 5–7, from cv. Colomba. Left line: molecular weight marker 100 bp (Gene Ruller, Fermentas).

confirmed by sequencing of the RT-PCR amplicon with primers P3/P4 (see Fig. 5, line 3). The partial sequence (204 bp) of this amplicon was identical (99.5 %) to PSTVd strain VP87 (LC523667) (Suppl. Fig. 2).


#### Stability of PSTVd in pure culture of *P. infestans*

The stability of strain VP87 in pure culture of *P. infestans* isolate PiVZR18v+ after consecutive passages on rye agar was studied. RT-PCR with primer sets P3/P4 (Fig. 6, a) and 68PV/87PV (Fig. 6, b) revealed amplification products



**Fig. 4.** RT-PCR detection of PSTVd with the primer set P3/P4 in *P. infestans* isolates grown on slices of potato cv. Colomba tubers after their isolation from infected with viroid tomato plants.

1, 2, Isolates from tomato cv. Moskvich infected by PSTVd strain VP87; 3, 4, isolates from tomato cv. Damskiy Palchik, infected by PSTVd strain VP87; 5–9, isolates from tomato cv. Moskvich, infected by PSTVd strain FP10-13; 10–13, isolates from tomato cv. Moskvich, infected by PSTVd strain VP35; 14, 15, isolates from tomato cv. Damskiy Palchik, infected by PSTVd strain VP35; 16, isolate PiVZR18 of *P. infestans* uninfected by PSTVd (negative control); 17, cv. Colomba potato tuber (negative control); 18, tomato cv. Moskvich infected by PSTVd strain VP87 (positive control); 19, distilled water (negative control). On the right and left sides of the gel, 50 bp DNA ladders (Primetech DNA Ladder) are shown.



**Fig. 5.** RT-PCR detection of PSTVd with the primer set P3/P4 in potato and tomato plants 60 dpi with viroid-infected *P. infestans* isolate PiVZR18v+.


1, Potato cv. Gala; 2–4, tomato cv. Zagadka; 5, tomato cv. Zagadka infected by PSTVd strain VP87 (positive control); 6, potato cv. Gala uninfected by PSTVd (negative control); 7, tomato cv. Zagadka uninfected by PSTVd (negative control); 8, PSTVd-uninfected *P. infestans* isolate PiVZR18 (negative control); 9, distilled water (negative control); M, molecular weight marker 100 bp (Gene Ruller, Fermentas).

indicating the presence of PSTVd after the second and third passages on rye agar of *P. infestans* isolates from viroid-infected cv. Colomba (see the Table, Fig. 6). PSTVd stability in *P. infestans* isolates after three passages on rye agar was shown by sequencing of the RT-PCR amplicon with primers P3/P4.

The partial sequence of RT-PCR amplicon (near 232 bp) of viroid in *P. infestans* isolate PiVZRv+ after the third passage on rye agar received with the primer set P3/P4 is identical (98.3 %) to PSTVd strain VP87 (LC523667). Another partial sequence of RT-PCR amplicon (270 bp) received with the primer set 68PV/87PV of the same RNA sample is identical to PSTVd strain VP87 (LC523667) – 99.3 % (Suppl. Fig. 3).

On the other hand, PSTVd was not detected after the eighth passage on rye agar in *P. infestans* subculture (see the Table).

*P. infestans* isolates infected with viroid strain VP87 were characterized by more abundant, but also more compact mycelium, forming an almost felt-like colony (Fig. 7).



**Fig. 6.** RT-PCR detection of PSTVd with primer sets P3/P4 (a) and 68PV/87PV (b) in cultures of *P. infestans* isolates after colonization on PSTVd infected potato cv. Colomba.


1–4, Second passage on rye agar; 5–11, third passage on rye agar; 12, PSTVd-uninfected isolate PiVZR18 (negative control); C<sup>+</sup>, cv. Colomba infected by PSTVd (positive control); C<sup>-</sup>, distilled water; M, molecular weight marker 100 bp (a) and 50 bp (b) (Gene Ruller, Fermentas).

Virulence testing on a set of Black's potato differentials (11 lines with different resistance genes) (Black et al., 1953) of the PSTVd-infected PiVZR18v+ and the initial uninfected PiVZR18 isolates showed the same types of reactions on detached leaves of 11 lines after 7 dpi (Fig. 8, 9).

Presence of amplification product 360 bp in *P. infestans* isolates after colonization on viroid-infected plants of potato cv. Colomba (each passage is 30 days)

| Number of passages on rye agar                     | Number of <i>P. infestans</i> isolates |                          |
|--|--|--------------------------|
|  | Positive PSTVd detection               | Negative PSTVd detection |
| One passage 30 days after re-isolation on rye agar | 5                                      | 2                        |
| Two passages                                       | 3 (see Fig. 6)                         | 1                        |
| Three passages                                     | 5 (see Fig. 6)                         | 2                        |
| Eight passages                                     | 0                                      | 7                        |


For both isolates, eight virulence genes (1, 2, 3, 4, 6, 7, 10, and 11) were identified. The PSTVd-infected *P. infestans* isolate seems to be less aggressive in comparison to the uninfected isolate; however, this observation requires further study.



**Fig. 7.** Cultures of *P. infestans* PiVZR18: a, infected with viroid after three passages on rye agar; b, healthy.



**Fig. 8.** Virulence of the PSTVd-infected PiVZR18v+ to the lines of a set of Black's differentials (11 resistance genes). Eight virulence genes (1, 2, 3, 4, 6, 7, 10, 11) were determined.



**Fig. 9.** Virulence of the initial PSTVd-uninfected *P. infestans* PiVZR18 isolate to the lines of a set of Black's differentials (11 resistance genes). Eight virulence genes (1, 2, 3, 4, 6, 7, 10, 11) were determined.

## Discussion

Potato (*Solanum tuberosum*) is one of the most important staple crops worldwide. According to the FAO, over 390 million tons are produced on over 19 million ha of farmland worldwide (<http://www.fao.org/faostat/en/#data/QCL>). The quarantine status of potato tuber spindle viroids and possible significant losses of potato yield determine the importance of studying various aspects of pathogen epidemiology.

PSTVd replication is accompanied by the accumulation of viroid-derived small RNAs (vd-siRNAs) suggested to play a central role in disease symptom development.

Potatoes were domesticated in the Andes in Southern Peru around 10,000 years ago. Nevertheless, the introduction of the potato to Europe, along with all its associated diseases, dates back to the 16–17th centuries (Khavkin, 2015). Late blight of potato and tomato is a devastating disease caused by fungus-like oomycete *P. infestans*. Despite many efforts, the severity of this disease has increased dramatically in recent years.

It is known that the mode and speed of spreading plant pathogens are the major factors in the development of epiphytotics. Early studies showed that PSTVd is spread primarily through the use of infected plant material produced vegetatively or as botanical seeds (Fernow et al., 1970), through mechanical spreading across the growing crop, particularly between plants of different species of the Solanaceae family (Manzer, Merriam, 1961; Verhoeven, Roenhorst, 2010). The success of mechanical transmission depends on infected host plant species or cultivars, as well as the frequency and severity of the disease and the temperature (Bulletin OEPP/EPPPO Bulletin, 2011). Importantly, transmission to potato and other test plants by aphids (*Myzus persicae*) was successful only if PSTVd RNA was encapsidated by potato leafroll virus (PLRV) particles (Salazar et al., 1995; Querci et al., 1997).

Some viruses are spread by vectors, which can include pathogenic fungi (Andika et al., 2017; Sutela et al., 2019), oomycetes (Mascia et al., 2019), and nematodes (Brown et al., 1989; Singh S. et al., 2020). Bidirectional transfer between *Fusarium graminearum* and tobacco plants of hop stunt viroid (HSVd) during infection was shown by Wei et al. (2019). However, *Nicotiana benthamiana* is not a natural host for either HSVd or *F. graminearum*. Given this fact (Serra et al., 2020), more evidence is needed to validate the possibility of viroid transmission by phytopathogenic fungi. We showed the presence of PSTVd in *P. infestans* isolates after colonization on plants of three potato cultivars infected with viroid. After three passages on rye agar (30 days each), RT-PCR analysis showed the presence of viroid in pure cultures of *P. infestans*. The partial RT-PCR amplicon sequence of viroid in *P. infestans* isolate PiVZRv+ after the third passage on rye agar is identical (98.3–99.3 %) to PSTVd strain VP87 that was used for the initial inoculation.

Sixty days after inoculation of healthy tomato plants with *P. infestans* isolate carrying PSTVd, RT-PCR revealed a 360 bp amplification product, indicating successful infection of the plants. This is the first report of horizontal transfer of potato viroid PSTVd between *P. infestans* and host plants.

Moreover, there is evidence that small RNAs (sRNAs, approximately 20–30 nt) can horizontally transfer from microbes to plants and spread silencing information toward the targeted genes (Han, Luan, 2015). Small RNAs were also found in

fungi (Wang, Dean, 2020) and fungal-like *Oomycota* (Jahan et al., 2015). In addition, sRNAs of 19–40 nt were found from *P. infestans* (Vetukuri et al., 2012). There are numerous reports of sRNA cross-transfer between plants and pathogens (Zeng et al., 2019; Wang, Dean, 2020). sRNAs can be transported within an organism through the inner side of the plasma membrane (symplast), or cell wall (apoplast) (Wang, Dean, 2020). It is suggested that sRNAs are translocated by extracellular vesicles (EVs) from *Arabidopsis* to *P. capsica* (Hou et al., 2019) and *B. cinerea* (Cai et al., 2018). Furthermore, it is possible that interaction between oomycete and potato involves not only sRNA exchange but also the movement of larger viroid RNA molecules from mycelium into a plant and *vice versa*.

PSTVd replicates in the nucleus, traffics long distances in the phloem, and moves cell-to-cell via plasmodesmata in plants (Takeda, Ding, 2009). After the third subculture, PSTVd was detected from *P. infestans*, suggesting that PSTVd can replicate in the nucleus and locate to non-septate hyphae of *P. infestans* (see the Table). On the other hand, after the eighth subculture, PSTVd accumulation in *P. infestans* was not detected by RT-PCR. The same results were obtained by Wei et al. (2019), in which PSTVd was eliminated from *Cryphonectria parasitica*, *Valsa mali*, and *Fusarium graminearum* after eight subcultures. This disappearance could be caused by a defense mechanism against viroid, namely, the RNA silencing system. Viroids are the target of the RNA silencing system and become elicitors of the host defense system via RNA silencing (Cottilli et al., 2019; Wei et al., 2020). Thus, PSTVd could have been degraded by the silencing system, resulting in the elimination of PSTVd from *P. infestans*.

*Phytophthora infestans* produce sporangia on the surface of potato leaves, and then zoospores, released from sporangia, form walled cysts on the plant surface (Mazumdar et al., 2021). The cysts germinate and extend a germ tube into the leaves and stems of the host plants. PSTVd transferred from *P. infestans* to plants, suggesting that PSTVd was present not only in mycelium but also in sporangia and zoospores. Mature sporangia were dispersed by wind or water (Leesutthiphonchai et al., 2018). Thus, there is a possibility that PSTVd can spread long distances via infected sporangia. In contrast, there is still no evidence of viroid infection in isolates of *P. infestans* from field populations and the possibility of viroid stability in the mycelium of *P. infestans* in tubers is unclear.

Concerning mycoviruses, there are two hypotheses of their origin: the first states that they are of an unknown but ancient origin and have coevolved along with their hosts, the second one suggests they have relatively recently moved from a fungal plant host into fungus (Pearson et al., 2009). Both hypotheses are also applicable to PSTVd. Prolonged coexistence of viroid–*P. infestans*–host plants can lead to viroid transition from a host plant to an oomycete.

## Conclusion

Potato spindle tuber viroid is known as autonomously replicating pathogen only of plants and mainly of solanaceous crops, that lacks any protein-coding sequences. Herein, we demonstrate the possibility of viroid transmission from host plants (potato and tomato) to *Phytophthora infestans*, from *P. infestans* to host plants, and the possibility of PSTVd

stability in pure cultures of *P. infestans* after three consecutive passages on rye agar. These results are initial evidence of bidirectionally transferred potato viroid PSTVd between *P. infestans* and host plants.

## References

- Andika I.B., Weia Sh., Cao Ch., Salaipetha L., Kondo H., Suna L. Phytopathogenic fungus hosts a plant virus: a naturally occurring cross-kingdom viral infection. *Proc. Natl. Acad. Sci. USA*. 2017;114(46):12267-12272. DOI 10.1073/pnas.1714916114.
- Behjatnia A., Dry I., Krake L., Condé B.D., Connelly M.I., Randles J., Rezaian M.A. New potato spindle tuber viroid and tomato leaf curl geminivirus strains from a wild *Solanum* sp. *Phytopathology*. 1996;86:880-886. DOI 10.1094/Phyto-86-880.
- Bhat A.L.I., Rao G.P. Characterization of Plant Viruses. Methods and Protocols. Humana, New York: Springer Science+Business Media, 2020. DOI 10.1007/978-1-0716-0334-5\_1.
- Black W., Mastenbroek C., Mills W.R., Peterson L.C. A proposal for an international nomenclature of races of *Phytophthora infestans* and of genes controlling immunity in *Solanum demissum* derivatives. *Euphytica*. 1953;2:173-178. DOI 10.1007/BF00053724.
- Brown D.J.F., Ploeg A.T., Robinson D.J. A review of reported associations between *Trichodorus* and *Paratrachodorus* species (Nematoda: Trichodoridae) and tobnaviruses with a description of laboratory methods for examining virus transmission by trichodorids. *Rev. Nematol.* 1989;12(3):235-241.
- Bulletin OEPP/EPPO Bulletin. 2011. *Potato spindle tuber viroid* on potato. 2011;41(3):394-399.
- Cai Q., Qiao L., Wang M., He B., Lin F.M., Palmquist J., Huang S.D., Jin H. Plants send small RNAs in extracellular vesicles to fungal pathogen to silence virulence genes. *Science*. 2018;360:1126-1129. DOI 10.1126/science.aar4142.
- Cottilli P., Belda-Palazón B., Adkar-Purushothama C.R., Perreault J.-P., Schleiff E., Rodrigo I., Ferrando A., Lisón P. Citrus exocortis viroid causes ribosomal stress in tomato plants. *Nucleic Acids Res.* 2019;47(16):8649-8661. DOI 10.1093/nar/gkz679.
- Diener T.O., Raymer W.B. Potato spindle tuber virus: a plant virus with properties of a free nucleic acid. II. Characterization and partial purification. *Virology*. 1969;37(3):351-366. DOI 10.1016/0042-6822(69)90219-0.
- Fernow K.H., Peterson L.C., Plaisted R.L. Spindle tuber virus in seeds and pollen of infected plants. *Am. Potato J.* 1970;47:75-80. DOI 10.1007/BF02864807.
- Gross H.J., Domdey H., Lossow C., Jank P.R.M., Alberty H., Sanger H.L. Nucleotide sequence and secondary structure of potato spindle tuber viroid. *Nature*. 1978;273:203-208. DOI 10.1038/273203a0.
- Han L., Luan Y.-S. Horizontal transfer of small RNAs to and from plants. *Front. Plant Sci.* 2015;6:1113. DOI 10.3389/fpls.2015.01113.
- Hou Y., Zhai Y., Feng L., Karimi H.Z., Rutter B.D., Zeng L., Choi D.S., Zhang B., Gu W., Chen X., Ye W., Innes R.W., Zhai J., Ma W. A *Phytophthora* effector suppresses trans-kingdom RNAi to promote disease susceptibility. *Cell Host Microbe*. 2019;25(1):153-165.e5. DOI 10.1016/j.chom.2018.11.007.
- Jahan S.N., asman A.K.M., Corcoran P., Fogelqvist J., Vetukuri R.R., Dixelius C. Plant-mediated gene silencing restricts growth of the potato late blight pathogen *Phytophthora infestans*. *J. Exp. Bot.* 2015;66:2785-2794. DOI 10.1093/jxb/erv094.
- Kastalyeva T.B., Mozhaeva K.A., Pisetskaya N.F., Romanova S.A., Trofimets L.N. The potato spindle tuber viroid and bringing potato into a healthy state. *Vestnik RASKHN = Herald of the Russian Academy of Agricultural Sciences*. 1992;3:22-24. (in Russian)
- Khavkin E.E. Potato late blight as a model of pathogen-host plant co-evolution. *Russ. J. Plant Physiol.* 2015;62(3):408-419. DOI 10.1134/S1021443715030103.
- Kryczyński S., Paduch-Cichal E.L., Skrzeczkowski J. Transmission of three viroids through seed and pollen of tomato plants. *J. Phytopathol.* 1988;121(1):51-57. DOI 10.1111/j.1439-0434.1988.tb00952.x.
- Leesuthiphonchai W., Vu A.L., Ah-Fong A.M.V., Judelson H.S. How does *Phytophthora infestans* evade control efforts? Modern insight into the late blight disease. *Phytopathology*. 2018;108(8):916-924. DOI 10.1094/PHYTO-04-18-0130-IA.
- Mackie A.E., Rodoni B.C., Barbetti M.J., McKirdy S.J., Jones R.A.C. Potato spindle tuber viroid: alternative host reservoirs and strain found in a remote subtropical irrigation area. *Eur. J. Plant Pathol.* 2016;145(2):433-446. DOI 10.1007/s10658-016-0857-2.
- Manzer F.E., Merriam D. Field transmission of potato spindle tuber virus and virus X by cultivating and hilling equipment. *Am. Potato J.* 1961;38:346-352.
- Mascia T., Vučurović A., Minutillo S.A., Nigro F., Labarile R., Savoia M.A., Palukaitis P., Gallitelli D. Infection of *Colletotrichum acutatum* and *Phytophthora infestans* by taxonomically different plant viruses. *Eur. J. Plant Pathol.* 2019;153(4):1001-1017. DOI 10.1007/s10658-018-01615-9.
- Matsushita Y., Yanagisawa H. Distribution of *Tomato planta macho viroid* in germinating pollen and transmitting tract. *Virus Genes*. 2018;54:124-129. DOI 10.1007/s11262-017-1510-7.
- Matsushita Y., Yanagisawa H., Khiutti A., Mironenko N., Ohto Y., Afanasenko O. Genetic diversity and pathogenicity of potato spindle tuber viroid and chrysanthemum stunt viroid isolates in Russia. *Eur. J. Plant Pathol.* 2021;161:529-542. DOI 10.1007/s10658-021-02339-z.
- Mazumdar P., Singh P., Kethiravan D., Ramathani I., Ramakrishnan N. Late blight in tomato: insights into the pathogenesis of the aggressive pathogen *Phytophthora infestans* and future research priorities. *Planta*. 2021;253(6):119. DOI 10.1007/s00425-021-03636-x.
- Medina M.V., Platt H.W. Comparison of different culture media on the mycelial growth, sporangia and oospore production of *Phytophthora infestans*. *Am. J. Potato Res.* 1999;76:121-125. DOI 10.1007/BF02853576.
- Mertelik J., Kloudova K., Cervena G., Necekalova J., Mikulkova H., Levkanicova Z., Dedic P., Ptacek J. First report of *Potato spindle tuber viroid* (PSTVd) in *Brugmansia* spp., *Solanum jasminoides*, *Solanum muricatum* and *Petunia* spp. in the Czech Republic. *Plant Pathol.* 2010;59(2):392. DOI 10.1111/j.1365-3059.2009.02115.x.
- Owens R.A., Khurana S.M.P., Smith D.R., Singh M.N., Garg I.D. A new mild strain of potato spindle tuber viroid isolated from wild *Solanum* spp. in India. *Plant Dis.* 1992;76(5):527-529. DOI 10.1094/PD-76-0527.
- Pearson M.N., Beever R.E., Boine B., Arthur K. Mycoviruses of filamentous fungi and their relevance to plant pathology. *Mol. Plant Pathol.* 2009;10(1):115-128. DOI 10.1111/j.1364-3703.2008.00503.x.
- Pfannenstiel M.A., Slack S.A. Response of potato cultivars to infection by the potato spindle tuber viroid. *Phytopathology*. 1980;70(9):922-926. DOI 10.1094/Phyto-70-922.
- Puchta H., Herold T., Verhoeven K., Roenhorst A., Ramm K., Schmidt-Puchta W., Sanger H.L. A new strain of potato spindle tuber viroid (PSTVd-N) exhibits major sequence differences as compared to all other PSTVd strains sequenced so far. *Plant Mol. Biol.* 1990;15(3):509-511. DOI 10.1007/BF00019169.
- Querci M., Owens R.A., Bartolini I., Lazarte V., Salazar L.F. Evidence for heterologous encapsidation of potato spindle tuber viroid in particles of potato leafroll virus. *J. Gen. Virol.* 1997;78(Pt.6):1207-1211. DOI 10.1099/0022-1317-78-6-1207.
- Ristaino J.B., Madritch M., Trout C.L., Parra G. PCR amplification of ribosomal DNA for species identification in the plant pathogen genus *Phytophthora*. *Appl. Environ. Microbiol.* 1998;64(3):948-954. DOI 10.1128/AEM.64.3.948-954.
- Salazar L.F., Querci M., Bartolini I., Lazarte V. Aphid transmission of potato spindle tuber viroid assisted by potato leafroll virus. *Fitopatologia*. 1995;30(1):56-58.
- Serra P., Carbonell A., Navarro B., Gago-Zachert S., Li S., Di Serio F., Flores R. Symptomatic plant viroid infections in phytopathogenic fungi: a request for a critical reassessment. *Proc. Natl. Acad. Sci. USA*. 2020;117(19):10126-10128. DOI 10.1073/pnas.1922249117.



- Singh R.P. Seed transmission of potato spindle tuber virus in tomato and potato. *Am. Potato J.* 1970;47:225-227. DOI 10.1007/BF02872303.
- Singh R.P. Experimental host range of the potato spindle tuber 'virus'. *Am. Potato J.* 1973;50:111-123. DOI 10.1007/BF02857207.
- Singh R.P., Boucher A., Somerville T.H. Detection of *Potato spindle tuber viroid* in the pollen and various parts of potato plant pollinated with viroid-infected pollen. *Plant Dis.* 1992;76:951-953.
- Singh S., Awasthi L.P., Jangre A., Nirmalkar V.K. Chapter 22 – Transmission of plant viruses through soil-inhabiting nematode vectors. In: Awasthi L.P. (Ed.) *Applied Plant Virology*. Acad. Press, 2020;291-300. DOI 10.1016/B978-0-12-818654-1.00022-0.
- Sutela S., Poimala A., Vainio E.J. Viruses of fungi and oomycetes in the soil environment. *FEMS Microbiol. Ecol.* 2019;95(9):fiz119. DOI 10.1093/femsec/fiz119.
- Syller J., Marczewski W., Pawłowicz J. Transmission by aphids of potato spindle tuber viroid encapsidated by potato leafroll luteovirus particles. *Eur. J. Plant Pathol.* 1997;103:285-289. DOI 10.1023/A:1008648822190.
- Takeda R., Ding B. Viroid intercellular trafficking: RNA motifs, cellular factors and broad impacts. *Viruses.* 2009;1(2):210-221. DOI 10.3390/v1020210.
- Verhoeven J.T.J., Roenhorst J.W. High stability of original predominant pospiviroid genotypes upon mechanical inoculation from ornamentals to potato and tomato. *Arch. Virol.* 2010;155:269-274. DOI 10.1007/s00705-009-0572-9.
- Vetukuri R.R., Åsman A.K.M., Tellgren-Roth C., Jahan S.N., Reimegård J., Fogelqvist J., Savenkov E., Söderbom F., Avrova A.O., Whisson S.C., Dixelius C. Evidence for small RNAs homologous to effector-encoding genes and transposable elements in the oomycete *Phytophthora infestans*. *PLoS One.* 2012;7(12):e51399. DOI 10.1371/journal.pone.0051399.
- Wang M., Dean R.A. Movement of small RNAs in and between plants and fungi. *Mol. Plant Pathol.* 2020;21:589-601. DOI 10.1111/mpp.12911.
- Wei Sh., Bian R., Andika I.B., Niu E., Liu Q., Kondo H., Yang L., Zhou H., Pang T., Lian Z., Liu X., Wu Y., Sun L. Symptomatic plant viroid infections in phytopathogenic fungi. *Proc. Natl. Acad. Sci. USA.* 2019;116(26):13042-13050. DOI 10.1073/pnas.1900762116.
- Wei S., Bian R., Andika I.B., Niu E., Liu Q., Kondo H., Yang L., Zhou H., Pang T., Lian Z., Liu X., Wu Y., Sun L. Nucleotide substitutions in plant viroid genomes that multiply in phytopathogenic fungi. *Proc. Natl. Acad. Sci. USA.* 2020;117(19):10129-10130. DOI 10.1073/pnas.2001670117.
- White T.J., Bruns T., Lee S., Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis M.A., Gelfand D.H., Sninsky J.J., White T.J. (Eds.) *PCR Protocols, a Guide to Methods and Applications*. Acad. Press, 1990;315-322. DOI 10.1016/B978-0-12-372180-8.50042-1.
- Yanagisawa H., Sano T., Hase S., Matsushita Y. Influence of the terminal left domain on horizontal and vertical transmissions of tomato planta macho viroid and potato spindle tuber viroid through pollen. *Virology.* 2019;526(2):22-31. DOI 10.1016/j.virol.2018.09.021.
- Zeng J., Gupta V.K., Jiang Y., Yang B., Gong L., Zhu H. Cross-kingdom small RNAs among animals, plants and microbes. *Cells.* 2019;8(4):371. DOI 10.3390/cells8040371.

---

#### ORCID ID

O.S. Afanasenko [orcid.org/0000-0001-7368-0797](https://orcid.org/0000-0001-7368-0797)  
A.V. Khiutti [orcid.org/0000-0003-1479-7746](https://orcid.org/0000-0003-1479-7746)  
N.V. Mironenko [orcid.org/0000-0003-3383-2973](https://orcid.org/0000-0003-3383-2973)  
N.M. Lashina [orcid.org/0000-0003-1419-7134](https://orcid.org/0000-0003-1419-7134)

**Acknowledgements.** This study was supported by the Russian Science Foundation, project No. 20-46-07001.

**Conflict of interest.** The authors declare no conflict of interest.

Received December 27, 2021. Revised February 3, 2022. Accepted February 3, 2022.