


doi 10.18699/vjgb-25-140

Overcoming the problem of heterologous proteins folding to improve the efficiency of yeast bioproducers

N.V. Dorogova , S.A. Fedorova 

Institute of Cytology and Genetics of the Siberian Branch of the Russian Academy of Sciences, Novosibirsk, Russia

 dorogova@bionet.nsc.ru

Abstract. In the last few decades, yeasts have been successfully engineered to be an excellent microbial cell factory for producing recombinant proteins with desired properties. This was due to their cost-effective characteristics and the successful application of genomic modification technologies. In addition, yeasts have a conserved post-translational modification pathway among eukaryotic organisms, which ensures the correct folding of recombinant proteins. However, the folding machinery cannot always cope with the load caused by the overexpression of recombinant genes, leading to the accumulation of misfolded proteins, the formation of aggregates and low production. Therefore, the protein-folding capacity of the endoplasmic reticulum (ER) remains one of the main limitations for heterologous protein production in yeast host organisms. However, thanks to many years of effective research of the fundamental mechanisms of protein folding, these limitations have been largely overcome. The study of folding in both model organisms and bioproducers has allowed to identify the molecular factors and cellular mechanisms that determine how a nascent polypeptide chain acquires its three-dimensional functional structure. This knowledge has become the basis for developing new effective techniques for engineering highly productive yeast strains. In this review, we examined the main cellular mechanisms associated with protein folding, such as ER transition, chaperone binding, oxidative folding, glycosylation, protein quality control. We discuss the effectiveness of applying this knowledge to the development of various engineering techniques aimed at overcoming bottlenecks in the protein folding system. In particular, selection of optimal signal peptides, co-expression with chaperones and foldases, modification of protein quality control, inhibition of proteolysis, and other techniques have allowed to enhance the ability of yeast bioproducers to effectively secrete heterologous proteins.

Key words: yeast bioproducers; protein folding; endoplasmic reticulum; molecular chaperones; recombinant proteins

For citation: Dorogova N.V., Fedorova S.A. Overcoming the problem of heterologous proteins folding to improve the efficiency of yeast bioproducers. *Vavilovskii Zhurnal Genetiki i Seleksii = Vavilov J Genet Breed.* 2025;29(8):1338-1347. doi 10.18699/vjgb-25-140

Funding. This study was supported by the Ministry of Science and Higher Education of the Russian Federation, project No. FWNR-2025-0031.

Решения проблем фолдинга белков для повышения эффективности дрожжевых биопродукторов

Н.В. Дорогова , С.А. Федорова 

Федеральный исследовательский центр Институт цитологии и генетики Сибирского отделения Российской академии наук, Новосибирск, Россия

 dorogova@bionet.nsc.ru

Аннотация. За последние несколько десятилетий дрожжи стали наиболее эффективными биопродукторами рекомбинантных белков с различными потребительскими свойствами. Это стало возможным благодаря их экономически выгодным характеристикам и успешному применению генно-инженерных технологий. Кроме того, дрожжи обладают консервативным для эукариотических организмов механизмом посттрансляционной модификации белков, который обеспечивает их корректный фолдинг, необходимый для дальнейшей секреции и функциональной активности. Однако аппарат фолдинга не всегда справляется с нагрузкой, вызванной сверхэкспрессией рекомбинантных генов, что приводит к накоплению неправильно свернутых белков, образованию агрегатов и низкой продуктивности дрожжевых штаммов. Таким образом, способность к фолдингу белков в эндоплазматическом ретикулеуме по-прежнему остается одним из основных ограничений при синтезе рекомбинантных белков в дрожжевых клетках. Эти ограничения были в значительной степени преодолены благодаря многолетним эффективным исследованиям фундаментальных механизмов белкового фолдинга. Изучение фолдинга как у модельных организмов, так и у биопродукторов позволило выявить молекулярные факторы и клеточные механизмы, определяющие формирование трехмерной функциональной структуры растущей пептидной цепи. Полученные знания легли в основу разработки новых эффективных методов конструирования высокопродуктивных штаммов дрожжей. В данном обзоре мы рассмотрели основные клеточные механизмы, связанные с фолдингом белков, такие как транспорт через эндоплазматический

ретикулум, взаимодействие с шаперонами, окислительный фолдинг, гликозилирование и контроль качества белков. Мы обсудили эффективность применения этих знаний при разработке различных инженерных методов, направленных на преодоление узких мест в системе белкового фолдинга. В частности, подбор оптимальных сигнальных пептидов, коэкспрессия с шаперонами и фолдазами, модификация клеточных механизмов контроля качества белков, ингибирование протеолиза и некоторые другие приемы позволили повысить возможности использования дрожжей-продуцентов в качестве эффективной производительной платформы для экспрессии и секреции рекомбинантных белков.

Ключевые слова: дрожжи-продуценты; фолдинг белков; эндоплазматический ретикулум; молекулярные шапероны; рекомбинантные белки

Introduction

Yeast expression systems are excellent for the production of valuable recombinant proteins and peptides widely used as biopharmaceuticals and industrial enzymes. They are commercially viable bioproducers due to their high growth rate, resistance to harmful microbiota, ability to assimilate many food sources, and fairly easy to cultivate in industrial conditions (Thak et al., 2020; Madhavan et al., 2021; De Brabander et al., 2023). The development of genetic and metabolic engineering has increased the efficiency of yeast strains, mainly due to the use of genomic technologies: strong promoters, new vector elements with improved inducers and enhancers, targeted mutagenesis, signaling molecules, high-performance devices for cloning, screening and fermentation (De Brabander et al., 2023; Tsuda, Nonaka, 2024). However, expression at the transcriptional and translational levels often does not correlate with the level of secretion of heterologous proteins, which is due to the insufficient efficiency of the folding mechanism (Ishiwata-Kimata, Kimata, 2023; Zahrl et al., 2023). Therefore, the search for new technological methods for optimizing the synthesis and increasing the yield of heterologous proteins in yeast cells remains an urgent task. A significant direction for its solution is overcoming the problem of folding target proteins into the correct three-dimensional structure.

Proper folding is necessary for the functional activity of synthesized proteins, their intracellular transport and further secretion. Recombinant proteins are secretory and go through a secretory pathway, beginning with folding in the ER and ending with release into the extracellular environment (culture media) (Raschmanová et al., 2021). Proteins are translocated to the ER in an unfolded state and then undergo modification and folding involving ER-resident chaperones, folding enzymes, and glycosylation (Hartl et al., 2011; Saibil, 2013). Disruptions in this machinery result in the accumulation of misfolded proteins in the cytoplasm, where they are recognized by the Protein Quality Control (PQC) system that regulates cellular homeostasis (Korennikh, Walter, 2012). This system includes the Unfolded Protein Response (UPR) signaling pathway, which can trigger refolding of misfolded proteins or initiate their proteolysis.

In some cases, misfolded proteins clump together to form aggregates or Inclusion Bodies (IBs), which can cause cell damage (Yamaguchi, Miyazaki, 2014). Such IBs often contain potentially active proteins with a normal secondary structure, which can be recovered from these aggregates under appropriate conditions (Burgess, 2009; Yamaguchi, Miyazaki, 2014; Singhvi et al., 2021). However, their isolation and refolding procedures are complex, expensive, and inefficient (Singhvi et al., 2021). It is therefore clear that the solution to the folding problem of heterologous proteins must be directly linked

to their production process: protein folding engineering and quality control in yeast host strains. In this article, we review various cellular mechanisms and signaling pathways that influence heterologous protein folding and discuss the latest updates to biotechnological strategy allowing to address this issue in order to maximize the yield of recombinant proteins.

Post-translational modifications and folding

Preparation for folding

In yeast, as in other eukaryotes, a newly synthesized peptide must undergo post-translational modification and folding in the endoplasmic reticulum to form the correct spatial conformation. A stable 3D structure determines the functional activity of the protein and its subsequent traffic through the secretory pathway (Schwarz, Blower, 2016).


Translation of secretory proteins occurs on cytosolic ribosomes and they are then co-translationally or post-translationally directed to the ER by a specific protein-RNA complex – Signal Recognition Particle (SRP). SRP binds to the N-terminal sequence of a precursor protein, which is called the signal peptide (SP). Transfer across the ER membrane occurs in an unfolded state of the nascent protein and is dependent on the Sec water channel, a multiprotein complex that spans the membrane (Berner et al., 2018; O’Keefe et al., 2022) (see the Figure).

When the unfolded nascent chain appears in the ER lumen, its hydrophobic sequence elements are recognized by ER-resident HSP chaperones: Kar2 (yeast Hsp70) and Ydj1 (yeast Hsp40) (Braakman, Hebert, 2013; Hendershot et al., 2024). They bind to the hydrophobic amino acid side chains exposed by the unfolded proteins. There is evidence that Kar2 assists in the folding of nascent proteins as they enter the ER and remains bound until folding is complete. The Ydj1 chaperone forms transient complexes with Kar2, facilitating its binding to non-native polypeptides (see the Figure). Thus, chaperones prevent premature misfolding of the immature polypeptide chain and protect it from aggregation (Omkar et al., 2024; Ruger-Herreros et al., 2024).

According to some data, the chaperone function of Kar2 in these events also depends on members of the DnaJ-like protein family, such as Jem1 and Scj1, and the nucleotide exchange factor Lhs1 (see the Figure). These co-chaperones promote the ATPase cycle and thus maintain Kar2 activity (Schlenstedt et al., 1995; Steel et al., 2004; de Keyzer et al., 2009).

Oxidative protein folding

3D structure formation begins with the process called Oxidative Protein Folding (OPF), which results in the formation of disulfide bonds due to the oxidation of thiol groups of cysteine



Nascent secretory protein is transferred co-translationally into the ER lumen through the Sec multiprotein complex that spans the ER membrane. At the ER membrane, the protein undergoes N-glycosylation involving the OST complex. The N-glycan-modified peptide chain binds to calnexin (Cnx). When the unfolded nascent chain appears in the ER lumen, it interacts with Kar2 and Ydj1. Jem1, Scj1 and Lhs1 assist in this process. In the ER lumen, the nascent peptide undergoes an OPF process involving Pdi and Ero, which results in disulfide bonds. Accumulation of misfolded proteins in the cytoplasm induces the UPR response which can trigger refolding of misfolded proteins or initiate ERAD.

(Hatahet, Ruddock, 2009; Palma et al., 2023). OPF is carried out by Protein disulfide isomerase (Pdi) (see the Figure). Pdi not only is responsible for the formation of the disulfide bonds in unfolded eukaryotic proteins, but also catalyzes the rearrangement of incorrect disulfide bonds (isomerase activity) (Gross et al., 2006). The OPF pathway, like Pdi, is conserved across eukaryotic organisms. Yeast Pdi1 protein is encoded by the *pdi1* gene, and *pdi1*-deleted yeast strains have low viability and accumulate secretory proteins within the ER lamellae (Mizunaga et al., 1990; Frand, Kaiser, 1998).

The transition of Pdi to the active form is catalyzed by ER sulfhydryl oxidase 1 (Ero1). Ero1 is an oxidoreductase that oxidizes Pdi1 via direct thiol-disulfide exchange (conversion of cystine to cysteine) (Gross et al., 2006; Sevier, Kaiser, 2006). Yeast contains a single *ero1* gene that encodes the Ero1 protein. The loss of *ero1* is lethal for yeast (Niu et al., 2016). Thus, Pdi and Ero act synergistically; for the correct catalysis of disulfide bonds in proteins, a balance of these two factors is necessary (Niu et al., 2016; Wang L., Wang C.C., 2023). The need to maintain such a balance is also due to the fact that the activation of Pdi and Ero occurs via the oxidation–reduction type and the high rate of disulfide bond formation in cells and tissues should create dangerous levels of oxidative stress (Gasser et al., 2008).

Role of glycosylation in promoting protein folding

Proper folding of most proteins requires post-translational modification known as glycosylation. In yeast, secretory proteins are glycoproteins and they contain covalently linked oligosaccharides, which are mainly mannose residues (see the Figure). Predominantly, yeast polypeptides are N-glycosylated, i.e. mannose is N-glycosidically linked to the β -amido group of asparagine. This reaction is catalyzed by the Oligosaccharyltransferase Enzyme Complex (OST) (Kelleher, Gilmore, 2006) and occurs on ER membranes. OST transfers Glc3Man9GlcNAc2-oligosaccharide (where Glc is glucose, Man is mannose, and GlcNAc is N-acetylglucosamine) from the lipid-pyrophosphate donor, dolichol diphosphate, to asparagine residues of nascent polypeptide chains. The protein-linked oligosaccharide is called N-glycan. This glycan has maintained a well-conserved structure throughout evolution and is characteristic of all eukaryotes (Qi et al., 2020). Glycans undergo further processing in the ER by glycosidases. In *Saccharomyces cerevisiae*, the following glycosidases have been described: α -glucosidase I encoded by CWH41, α -glucosidase II encoded by ROT2, α 1,2-mannosidase encoded by MNS1. Glycosidases partially deglycosylate and shorten the N-glycan (Herscovics, 1999; Lehle et al., 2006).

Glycosylation promotes folding by enhancing the solubility and stability of the proteins in the ER and the Golgi. Oligosaccharide residues are a marker for interactions with certain chaperones, also assisting in the folding of glycoproteins (Parodi, 2000; Xu C., Ng, 2015).

The N-glycan-modified polypeptide chain is recognized by lectin chaperones (Ware et al., 1995; Caramelo, Parodi, 2015). In mammalian cells, two related ER lectin chaperones, calnexin (Cnx) and calreticulin (Crt), are important for the proper folding of newly synthesized glycoproteins. Calnexin is an integral membrane protein, and calreticulin is a soluble protein found in the ER lumen (see the Figure). They retain glycoproteins in the ER during translation by inhibiting their aggregation and formation of non-canonical disulfide bridges, and also promote their association with other chaperones. In yeast (*S. cerevisiae*), only the calnexin homologue Cne1 has been identified. Cne1 is structurally similar to mammalian calnexin except that it lacks a cytoplasmic tail and does not bind calcium (Parlati et al., 1995). Yeast calnexin has been shown to function as a molecular chaperone similar to mammalian calnexin (Xu X. et al., 2004). A calnexin-like transmembrane protein has also been identified in *S. pombe* (Núñez et al., 2015). However, genes encoding a calreticulin homologue have not yet been identified in the yeast genome.

Protein Quality Control

All eukaryotes have conserved mechanisms that control cellular proteostasis and protect the cell from stress. These are the three main pathways that comprise the quality control system: UPR, ERAD (ER-associated degradation), and autophagy. There are essentially two alternative cellular responses to the accumulation of abnormal proteins. UPR promotes their re-folding repair through additional activation of ER chaperones and folding enzymes, while ERAD and autophagy target them for degradation (see the Figure).

Unfolded protein response

In eukaryotes, UPR includes ER-localized molecular chaperones that participate in sensing misfolded proteins, activating downstream signaling cascades, and mitigating proteotoxic ER stress: Inositol-requiring enzyme 1 (Ire1), Activating transcription factor 6 (Atf6), and Protein kinase R-like ER kinase (PERK). However, only Ire1 is highly conserved and has been found in unicellular eukaryotes, including yeast (Schroder et al., 2003; Mori K., 2022). Presumably, through its luminal domain, Ire1 enables direct interaction with exposed hydrophobic groups of misfolded proteins in the ER. This reaction results in the activation of Ire1, and as a result, its C-terminal RNase domain mediates splicing of the *Hac1* gene mRNA removing a 252-nucleotide intron near the 3' end. Then Rlg1, a tRNA ligase, ligates the *hac1* transcript cleaved by Ire1p. Its spliced and unspliced forms are termed *Hac1i* and *Hac1u*, respectively ("i" and "u" for induced and uninduced) (Schroder et al., 2003; Xia, 2019). *Hac1i* mRNA is translated into the transcription factor Hac1. It is then transported into the nucleus where it induces transcription of a large number of genes involved in the UPR mechanism (Hernández-Elvira et al., 2018) including those encoding ER-located molecular chaperones and protein modification enzymes such as *kar2*, *pdi1*, *ero1*, *ecj1*, *lhs1*, *jem1*. Additionally, the Ire1/Hac1 pathway is essential for activating genes that carry out ERAD functions promoting

selective removal of terminally damaged proteins (Friedlander et al., 2000; Travers et al., 2000).

Hac1 orthologs have been identified in various yeast species, such as *Pichia pastoris*, *Hansenula polymorpha*, *Kluyveromyces lactis*, *Yarrowia lipolytica*, *Candida albicans*, and *Candida parapsilosis*. All of these species share the Ire1-dependent mechanism of splicing the *Hac1* transcripts in response to ER stress (Hernández-Elvira et al., 2018; Fauzee et al., 2020; Ishiwata-Kimata, Kimata, 2023).

However, it has been reported that *S. pombe* probably does not contain a *Hac1* ortholog. Its Ire1 triggers a process called Regulated Ire-dependent decay (Kimmig et al., 2012). In *S. pombe*, stress-activated IRE1 cleaves mRNAs located in the ER, leading to their exonuclease-mediated degradation (Hernández-Elvira et al., 2018).

ER-associated degradation

Proteins that have not achieved their native conformation after repeated folding remain in the ER, bound to ER chaperones that prevent their aggregation. If a protein remains unfolded in the ER for too long, it is identified as potentially harmful and eliminated via ER-associated degradation. The process involves several steps: the recognition of substrates in the lumen and membrane of the ER, their translocation into the cytosol, ubiquitination and degradation in the 26S proteasome. Thus, ERAD is associated with the highly conserved ubiquitin proteasome system (UPS) (Ruggiano et al., 2014; Krshnan et al., 2022). In all eukaryotes, it includes the following main components: ubiquitin-activating enzyme (E1), ubiquitin-conjugating enzyme (E2), ubiquitin ligase (E3), and 26S proteasome (Pickart, 2001). If a misfolded glycoprotein stays in the ER for a critically long time, its N-linked glycans are trimmed by yeast 1,2-mannosidase1 – Htm1. This trimmed N-glycan is recognized by a lectin chaperone known as Yos9. At the same time, the Hrd3 protein (HMG-CoA reductase degradation protein 3) associates with hydrophobic amino acid residues exposed on the surface of misfolded glycoproteins (Thibault, Ng, 2012; Berner et al., 2018). After recognition and binding to Hrd3 and Yos9, the substrate protein is transferred to the E3 ubiquitin ligases complex responsible for ubiquitination – Hrd1 (HMG-CoA reductase degradation protein 1). Hrd1 is embedded in the ER membrane and is in conjunction with Der1 (Degradation in the endoplasmic reticulum protein 1) and Usa 1 (U1 SNP1-associating protein 1), which mediates retrotranslocation of misfolded proteins through the ER membrane from the lumen side of the cytosolic membrane. The Hrd1 E3 ubiquitin ligase contains a RING domain that accepts ubiquitin from the membrane-associated protein Ubc7 (E2 ubiquitin-conjugating ligase). Finally, Hrd1 transfers ubiquitin to the substrate protein. Proteins covalently linked to one or more ubiquitin molecules are recognized by proteasome (Preston, Brodsky, 2017; Berner et al., 2018; Krshnan et al., 2022).

Ways to solve the protein folding problem in yeast synthetic biology

Selection of ER-targeting signal peptides

ER-targeting signal peptides (SPs) are critical components for the secretion of heterologous proteins because they are required for their correct transport and localization to the ER,

where modification and folding occur (Zha et al., 2023). SPs are short sequences, 15 to 30 amino acids in length, mostly located at the N-terminus of the secreted proteins. SPs typically consist of a series of hydrophobic amino acids (core region) that embed into membranes, positively charged residues at the N-terminus (necessary for proper topology of the polypeptide), and a cleavage site at the C-terminus. When an SP is cleaved by a signal peptidase, it releases the secreted protein into the ER lumen (Zha et al., 2023).

Recent research has revealed that the secretion efficiency of heterologous proteins is strongly dependent on their successful combination with SPs. Therefore, various bioinformatic and experimental studies have been carried out to elucidate the optimal SP sequence, allowing to maximize the efficiency of protein secretion.

A wide range of signal sequences, both from native genes (including those from different organisms) and synthetic molecules, are used to improve heterologous expression in yeast. For example, in *Komagataella phaffii* (*P. pastoris*), a widespread host microorganism, the most commonly used secretion signal is the mating pheromone of the α -factor (MF α 1) from *S. cerevisiae* (Eskandari et al., 2023). Although this SP has proven its effectiveness in increasing the yield of heterologous proteins, work continues on its modification and the search for more productive variants. In particular, it was shown that some single amino acid substitutions of the MF α signal sequences provided a significant increase in the production of secreted proteins (Ito et al., 2022). An improved secretion signal was also obtained by creating a chimeric construct combining the MF α 1 leader region and the Ost1 signal sequence. This hybrid variant turned out to be more effective compared to the original MF α 1 SP (Barrero et al., 2018). The emergence of new improved modifications of MF α 1 SPs was facilitated by the analysis of mutations accumulated in MF α signal sequences during yeast evolution. This allowed to identify specific motifs as well as their combinations (mutation synergism) that are important for enhancing yeast enzyme secretion (Aza et al., 2021). Promising results were obtained by combining bioinformatic prediction of the efficiency of certain signal sequences and subsequent experimental validation. For example, Duan and colleagues (2019) discovered four new endogenous signal peptides, including Dan4, Gas1, Msb2, and Fre2, according to the reported secretome and genome of *P. pastoris*. Their properties were investigated experimentally using three reporter proteins, and these SPs were shown to be superior to α -MFs in the production of heterologous proteins (Duan et al., 2019).

The SignalP databases (www.signalpeptide.de) can be used as a resource for selecting suitable SPs. Mori and colleagues (2015) created a library of 60 *S. cerevisiae* SPs that were identified in SignalP 3.0 using SOSUI software. The authors experimentally showed that six of those SPs can maximize secretion of heterologous proteins (Mori et al., 2015).

While some progress has been made in bioinformatically predicting the efficiency of particular signal peptides for recombinant proteins, their potential has only been explored in laboratory settings. There is no reliable information yet on their successful application in biotech platforms, and it is unclear how they will function in specific yeast strains and in combination with specific target proteins.

Increased activity of the ER folding network

Positive effects of chaperone addition

Traditionally (since the 90s), yeast strain engineering has used the method of simultaneous expression (co-expression) of genes encoding heterologous proteins and genes encoding chaperones and folding enzymes. It has been shown for a variety of heterologous proteins that the introduction of extra copies of these genes into yeast host cells increases the secretion and decreases the aggregation of recombinant proteins. For example, Robinson et al. (1994) demonstrated that overexpression of Pdi in *S. cerevisiae* led to a four- to tenfold increase in secretion yields of human protein. Shusta and co-authors (1998) reported 2–6-fold increased secretion titers for single-chain antibody fragments upon co-overexpression of the Kar2 chaperone or Pdi. Simultaneous overexpression of Pdi and Kar2 resulted in a synergistic up to eightfold increase (Shusta et al., 1998).

The addition of Pdi and Kar2, both together and separately, increased the yield of recombinant proteins used for medical purposes: antithrombotic factor, hirudin (Kim et al., 2003); mammalian peptide recognition proteins (Yang et al., 2016); *Necator americanus* secretory protein (Inan et al., 2007); fragment of single-chain antibody A33 (Damasceno et al., 2007); hydrophobins (Sallada et al., 2019), virus glycoprotein, RABV-G (Ben Azoun et al., 2016).

In some cases, chaperone activity is enhanced by their co-expression with molecular partners and cofactors. For example, Kar2 functionality depends on the ATPase cycle, which is promoted by the Jem1 co-chaperone and the Lhs1p nucleotide exchange factor (Steel et al., 2004). Joint expression of genes encoding these factors has been shown to stimulate Kar2 activity and increase production of recombinant human proteins (Payne et al., 2008).

As we wrote above, yeast strains overexpressing Pdi are the most frequently used in the production of correctly folded and functionally secreted recombinant proteins. However, in the oxidative protein folding pathway, Pdi acts in partnership with Ero1-oxidase. Therefore, co-overexpression of these genes can improve the efficiency of protein folding and secretion. Beal and colleagues (2019) developed a new methodology enabling the quantitative assessment of the interaction of Pdi1 and Ero1, and based on it provided a platform for the design of more efficient heterologous protein expression systems in yeast. The high efficiency of co-expression of Pdi and its molecular partner Ero1 was also shown by other authors (Ben Azoun et al., 2016; Sallada et al., 2019).

Thus, addition of known ER network folding factors to enhance heterologous expression has become a basic approach. However, this is not a universal solution; there is no guarantee that chaperones are always able to improve the secretion of recombinant proteins. In particular, a combination of different chaperones does not always lead to a synergistic effect.

Limitations of the approach

Recent investigations suggest that the efficiency of chaperones depends on the dose of the gene encoding the recombinant protein. For example, overexpression of Pdi1 or Ero1 caused a statistically significant increase in recombinant hydrophobin in *P. pastoris*, but only in the 3-copy gene strain, and

not in the 1- and 2-copy strains (Sallada et al., 2019). In the same experiment, it was shown that hydrophobin production increased with co-expression of Kar2 14-fold for the 1-copy strain, 9.8-fold for the 2-copy and 22-fold for the 3-copy ones (Sallada et al., 2019).

A correlation between chaperone activity and the copy number of the recombinant gene has been found in other experiments as well. In lipase-producing *P. pastoris*, overexpression of Pdi1 led to enhanced productivity in the strain carrying four copies of the lipase gene, whereas in the two-copy strain, it remained unchanged (Huang J. et al., 2020). Efficient expression of mammalian peptidoglycan recognition proteins in *P. pastoris* also depended on the combination of recombinant gene copy number and folding enzymes (Yang et al., 2016). Thus, these experiments support the need for empirical selection of optimal combinations of recombinant protein-encoding genes and folding factor-encoding genes.

Recently, an increasing number of published data have demonstrated that the activity of the folding factors described above is selective for certain substrates and does not improve the expression of some important recombinant proteins. In particular, it was shown that Pdi1, Ero1, Kar2 did not have a beneficial effect on the secretion of antibodies in *S. cerevisiae* (de Ruijter et al., 2016). Similar conclusions were also drawn from other experiments (Smith et al., 2004; Payne et al., 2008). These findings demonstrate that there is no single suitable strategy that provides optimal conditions for the secretion of all recombinant proteins, and very often a separate productivity enhancement program must be set up for each protein.

Forced activation of the UPR system

An effective way to solve the folding problem is the activation of the ER chaperone network by overexpression of the Hac1 transcription factor, which is the main regulating UPR pathway. Theoretically, an artificial increase in the level of Hac1 can be achieved in two ways. The first is the intensification of Hac1 mRNA splicing by overexpression of Ire1. The second way is overexpression of genetic sequences encoding Hac1.

The first approach is associated with additional expression of Ire1, which should lead to an increase in the number of events associated with Hac1 mRNA splicing and, as a consequence, promote transcription of genes encoding ER chaperones and folding catalysts (see above “Unfolded protein response”). However, at present, the effectiveness of this approach is confirmed by only one experiment. Only one study convincingly showed that overexpression of Ire1 improved the capabilities of the yeast expression system, which in particular led to an increase in hepatitis B small antigen (HBsAg) production in *S. cerevisiae* (Sheng et al., 2017). The lack of positive results may be due to the complex mechanism of Hac1 activation, which involves many molecular factors.

The second approach aimed at using Hac1 overexpression is the most preferable in yeast biotechnology. As we wrote above, adding Hac1 to yeast host strains leads to up-regulation of a large number of its target genes that mainly function for the protein secretory machinery, and in particular, almost all these genes are required for protein folding (see above “Unfolded protein response”). For heterologous protein secretion, both the active (spliced) form and the inactive (unspliced) form of

Hac1 can be used. However, activation of the unspliced form of Hac1 mRNA requires additional cellular resources, in particular, molecular factors involved in its processing. The genes encoding these factors must be added to the yeast strains. The engineered strains with such a modification increased protein secretion significantly and showed better performance than strains with only overexpression of Hac1 (Lin et al., 2023).

Most strains used in biotechnological practice contain an insertion of the Hac1 gene with a previously removed 3' end intron, i.e. encoding the active form of the transcript. For such strains, a high level of transcriptional activity of Hac1 target genes involved in the UPR pathway and responsible for correct protein folding was shown. In *P. pastoris* strains overexpressing Hac1, electron microscopy revealed an expansion of the intracellular membranes (Guerfal et al., 2010), which probably indicates an increase in the folding capacity of the ER and may contribute to the alleviation of ER stress (Schuck et al., 2009).

To date, a large number of research groups have reported an increase in the productivity of secretory proteins due to artificial and high-level expression of the Hac1 protein (Raschmanová et al., 2021; Lin et al., 2023; Khlebodarova et al., 2024). Hac1 overexpression successfully increased the yield of recombinant proteins: phytase (a product of the *Phy* gene from *C. amalonaticus*) in *P. pastoris* (Li C. et al., 2015); α -amylase (Valkonen et al., 2003; Lin et al., 2023); chitosanase (from *Bacillus subtilis*) (Han et al., 2021); krigle fragment of human apolipoprotein (which inhibits endothelial cell migration) in *S. cerevisiae* (Lee et al., 2012); xylanase (a microbial hydrolase used to hydrolyze xylan) in *S. cerevisiae* (Li C. et al., 2015; Bao et al., 2020).

However, to achieve efficient heterologous expression in Hac-modified cells, it is important to select the optimal combination of the copy number of Hac-encoding genes and genes encoding recombinant proteins (Valkonen et al., 2003; Guerfal et al., 2010; Huang M. et al., 2017; Huang J. et al., 2020). Some studies have found that increasing Hac1 doses also leads to increased ER stress (Gasser et al., 2006; Guerfal et al., 2010; Li C. et al., 2015). It is also necessary to take into account that Hac1 is a positive regulator of the ERAD pathway (see above “ER-associated degradation”) and its overactivation can promote protein degradation.

Thus, the addition of Hac allows the yeast strain to be adapted to large-scale protein expression caused by an excessive dose of transgenic constructs. However, many researchers note that the effectiveness of this approach must be assessed in each specific case when designing a specific producer strain.

An additional advantage in strain engineering may be gained by using Hac orthologs from other yeasts, and even phylogenetically more distant eukaryotes. In particular, Valkonen et al. (2003) reported that the secretion of α -amylase was increased by overexpressing *Trichoderma reesei*-derived Hac1 in *S. cerevisiae*. Bankefa and colleagues (2018) showed that for *P. pastoris*, Hac1 orthologs of other species and even mammalian ones may be more effective than the native one. They investigate the effects of overexpressing Hac1 orthologs from *S. cerevisiae* (ScHac1p), *Trichoderma reesei* (TrHac1p) and *Homo sapiens* (HsXbp1) on the secretory expression levels of three reporter proteins, b-galactosidase, b-mannanase and glucose oxidase. The authors reported diverse effects of

these orthologs on heterologous expression levels, but HsXbp1 remarkably improved the enzyme production levels, both in shake flask and fermenter culture, both in single- and four-copy strains, which demonstrated its great application potential (Bankefa et al., 2018).

Thus, artificial activation of the UPR pathway by overexpression of the Hac1 transcription factor has demonstrated an obvious positive effect on improving the secretory protein productivity. It allowed to remove bottlenecks in the engineered yeast strain, arising due to abnormal accumulation of unfolded/misfolded proteins in the endoplasmic reticulum. However, this positive experience cannot be extrapolated to all recombinant proteins. Therefore, the real effect for each product must be assessed in experiments, often based on trial and error.

Prevention of ERAD pathway activation

As we wrote above, the UPR system is in crosstalk with the ERAD pathway. ERAD is activated when ER chaperones and folding enzymes are unable to form tertiary or quaternary structures of proteins. Sometimes ERAD is excessively activated during heterologous expression, so depletion of some components of this system can contribute to an increase in the yield of recombinant proteins.

De Ruijter and Frey (2015) analyzed the effect of deletions of genes involved in ERAD on the production of human IgG in *S. cerevisiae*. It was shown that deletion of only one gene, HTM1, contributed to a slight improvement in heterologous secretion, whereas deletions of the *yos9*, *hrd1*, *hrd3*, and *ubc7* genes either did not affect or negatively affected the recombinant protein yield (de Ruijter, Frey, 2015).

In *P. pastoris*, excessive activation of ERAD enhanced intracellular degradation of recombinant antibody fragment Fab. In the work of Pfeffer et al., it was shown that most of the newly synthesized Fab is not secreted but undergoes intracellular degradation via the ubiquitin-proteasome system (Pfeffer et al., 2012; Zahrl et al., 2019). The yield of the recombinant protein was increased by inhibiting proteasome components (Pfeffer et al., 2012). However, subsequent work aimed at reducing proteolysis through ERAD gene disruptions did not yield significant increases in Fab secretion (Zahrl et al., 2019).

Thus, the reduction of ERAD activity can be considered as a potential strategy for improving the secretion of recombinant proteins. However, the current level of research in this area does not yet allow for the transition to engineering producer strains protected from the proteolysis system.

Search for new solutions

The potential of the ER folding network can be enhanced by factors that are not directly involved in it, but create favorable conditions for its active functioning.

Zahrl et al. (2023) proposed to combine the transcriptional programs induced by Hac1 and Msn4 in one strain. Msn4 is a transcriptional factor involved in the response to various forms of stress (heat, oxidative, osmotic, etc.). Co-expression of Hac1 and Msn4 (both native and synthetic) revealed synergistic effects resulting in increased titers of recombinant proteins. This strategy was tested for scFv and VHH antibody fragments expressed in *P. pastoris* (Zahrl et al., 2023).

In another work, the same research group identified the most relevant chaperones of the Hsp70 network, both cytosolic and ER-localized, and investigated the impact of their combined overexpression on recombinant protein secretion (Zahrl et al., 2022). In their work, they implemented a principle they called the push-and-pull strategy. The addition of cytosolic chaperones allowed to increase the translocation competency of the recombinant protein and its targeting to the ER membrane (= push). At the same time resident ER chaperones improved the folding process (= pull). This allowed to successfully engineer strains and improve protein secretion up to 5-fold for the antibody Fab fragment and scFv (Zahrl et al., 2022).

The screening of new molecules involved in the folding system may improve yeast expression systems and expand the range of heterologous proteins. In particular, analysis of the reported *P. pastoris* secretome and genome predicted novel folding factors: Mpd1 and Pdi2 (members of the Pdi family), as well as Sil1 (nucleotide exchange factor for Kar2) (Duan et al., 2019). Subsequent experimental studies showed that all of the novel folding factors enhanced total production of reporter proteins, with Sil1 showing the highest efficiency (Duan et al., 2019). This work is an example of a successful combination of the achievements of yeast omics technologies and metabolic engineering, but this experience has not yet been widely applied.

Conclusion

One of the main limitations of heterologous protein production in yeast hosts is the ability of proteins to fold in the endoplasmic reticulum. The folding system is subject to unbalanced stress due to overexpression of recombinant genes, leading to the accumulation of misfolded proteins, aggregate formation, and low productivity. However, thanks to years of effective research into the fundamental mechanisms of protein folding, these limitations have been largely overcome. Studying folding in both model organisms and bioproducers has enabled the identification of molecular factors and cellular mechanisms that determine how a nascent polypeptide chain acquires its three-dimensional functional structure. This knowledge has formed the basis for the development of new, efficient methods for constructing highly productive yeast strains. Many problems arising from insufficient folding systems have been overcome by selecting optimal signal peptides, coexpressing with chaperones and foldases, modifying the ubiquitin-proteasome system (UPS), and preventing the ERAD pathway. Modern engineering solutions utilize combinations of these factors, but for each protein of interest, the expression strain is typically developed individually. In practice, optimized folding conditions for one protein often do not work for another. Therefore, no general strategy for overcoming protein folding bottlenecks that would be applicable to a wide range of proteins has yet been proposed.

In the future, some problems can be minimized by analyzing data obtained using omics technologies and modeling the secretion pathway *in silico*. An example of such a development is the pcSecYeast model designed for *S. cerevisiae* (Li F. et al., 2022). Such models allow choosing a combination of factors, both known and unknown, to generate new engineering strategies in designing strains with high protein yields.

References

- Aza P., Molpeceres G., de Salas F., Camarero S. Design of an improved universal signal peptide based on the α -factor mating secretion signal for enzyme production in yeast. *Cell Mol Life Sci.* 2021;78(7):3691-3707. doi 10.1007/s00018-021-03793-y
- Bankefa O.E., Wang M., Zhu T., Li Y. Hac1p homologues from higher eukaryotes can improve the secretion of heterologous proteins in the yeast *Pichia pastoris*. *Biotechnol Lett.* 2018;40(7):1149-1156. doi 10.1007/s10529-018-2571-y
- Bao C., Li J., Chen H., Sun Y., Wang G., Chen G., Zhang S. Expression and function of an *Hac1*-regulated multi-copy xylanase gene in *Saccharomyces cerevisiae*. *Sci Rep.* 2020;10(1):11686. doi 10.1038/s41598-020-68570-6
- Barrero J.J., Casler J.C., Valero F., Ferrer P., Glick B.S. An improved secretion signal enhances the secretion of model proteins from *Pichia pastoris*. *Microb Cell Fact.* 2018;17(1):161. doi 10.1186/s12934-018-1009-5
- Beal D.M., Bastow E.L., Staniforth G.L., von der Haar T., Freedman R.B., Tuite M.F. Quantitative analyses of the yeast oxidative protein folding pathway *in vitro* and *in vivo*. *Antioxid Redox Signal.* 2019;31(4):261-274. doi 10.1089/ars.2018.7615
- Ben Azoun S., Belhaj A.E., Göngrich R., Gasser B., Kallel H. Molecular optimization of rabies virus glycoprotein expression in *Pichia pastoris*. *Microb Biotechnol.* 2016;9(3):355-368. doi 10.1111/1751-7915.12350
- Berner N., Reutter K.R., Wolf D.H. Protein quality control of the endoplasmic reticulum and ubiquitin-proteasome-triggered degradation of aberrant proteins: yeast pioneers the path. *Annu Rev Biochem.* 2018;20(87):751-782. doi 10.1146/annurev-biochem-062917-012749
- Braakman I., Hebert D.N. Protein folding in the endoplasmic reticulum. *Cold Spring Harb Perspect Biol.* 2013;5(5):a013201. doi 10.1101/cshperspect.a013201
- Burgess R.R. Refolding solubilized inclusion body proteins. *Methods Enzymol.* 2009;463:259-282. doi 10.1016/S0076-6879(09)63017-2
- Caramelo J.J., Parodi A.J. A sweet code for glycoprotein folding. *FEBS Lett.* 2015;589(22):3379-3387. doi 10.1016/j.febslet.2015.07.021
- Damasceno L.M., Anderson K.A., Ritter G., Cregg J.M., Old L.J., Batt C.A. Cooverexpression of chaperones for enhanced secretion of a single-chain antibody fragment in *Pichia pastoris*. *Appl Microbiol Biotechnol.* 2007;74(2):381-389. doi 10.1007/s00253-006-0652-7
- De Brabander P., Uitterhaegen E., Delmulle T., De Winter K., Soetaert W. Challenges and progress towards industrial recombinant protein production in yeasts: a review. *Biotechnol Adv.* 2023;64:108121. doi 10.1016/j.biotechadv.2023.108121
- de Keyser J., Steel G.J., Hale S.J., Humphries D., Stirling C.J. Nucleotide binding by Lhs1p is essential for its nucleotide exchange activity and for function *in vivo*. *J Biol Chem.* 2009;284(46):31564-31571. doi 10.1074/jbc.M109.055160
- de Ruijter J.C., Frey A.D. Analysis of antibody production in *Saccharomyces cerevisiae*: effects of ER protein quality control disruption. *Appl Microbiol Biotechnol.* 2015;99:9061-9071. doi 10.1007/s00253-015-6807-7
- de Ruijter J.C., Koskela E.V., Frey A.D. Enhancing antibody folding and secretion by tailoring the *Saccharomyces cerevisiae* endoplasmic reticulum. *Microb Cell Fact.* 2016;23(15):87. doi 10.1186/s12934-016-0488-5
- Duan G., Ding L., Wei D., Zhou H., Chu J., Zhang S., Qian J. Screening endogenous signal peptides and protein folding factors to promote the secretory expression of heterologous proteins in *Pichia pastoris*. *J Biotechnol.* 2019;306:193-202. doi 10.1016/j.jbiotec.2019.06.297
- Eskandari A., Nezhad N.G., Leow T.C., Rahman M.B.A., Oslan S.N. Current achievements, strategies, obstacles, and overcoming the challenges of the protein engineering in *Pichia pastoris* expression system. *World J Microbiol Biotechnol.* 2023;40(1):39. doi 10.1007/s11274-023-03851-6
- Fauzee Y.N.B.M., Taniguchi N., Ishiwata-Kimata Y., Takagi H., Kimata Y. The unfolded protein response in *Pichia pastoris* without external stressing stimuli. *FEMS Yeast Res.* 2020;20(7):foaa053. doi 10.1093/femsyr/foaa053
- Fränd A.R., Kaiser C.A. The *Ero1* gene of yeast is required for oxidation of protein dithiols in the endoplasmic reticulum. *Mol Cell.* 1998;1(2):161-170. doi 10.1016/S1097-2765(00)80017-9
- Friedlander R., Jarosch E., Urban J., Volkwein C., Sommer T. A regulatory link between ER-associated protein degradation and the unfolded-protein response. *Nat Cell Biol.* 2000;2(7):379-384. doi 10.1038/35017001
- Gasser B., Maurer M., Gach J., Kunert R., Mattanovich D. Engineering of *Pichia pastoris* for improved production of antibody fragments. *Biotechnol Bioeng.* 2006;94(2):353-361. doi 10.1002/bit.20851
- Gasser B., Saloheimo M., Rinas U., Dragosits M., Rodríguez-Carmona E., Baumann K., Giuliani M., ... Porro D., Ferrer P., Tutiño M.L., Mattanovich D., Villaverde A. Protein folding and conformational stress in microbial cells producing recombinant proteins: a host comparative overview. *Microb Cell Fact.* 2008;4(7):11. doi 10.1186/1475-2859-7-11
- Gross E., Sevier C.S., Heldman N., Vitu E., Bentzur M., Kaiser C.A., Thorpe C., Fass D. Generating disulfides enzymatically: reaction products and electron acceptors of the endoplasmic reticulum thiol oxidase Ero1p. *Proc Natl Acad Sci USA.* 2006;103(2):299-304. doi 10.1073/pnas.0506448103
- Guerfal M., Ryckaert S., Jacobs P.P., Ameloot P., Van Craenenbroeck K., Derycke R., Callewaert N. The *HAC1* gene from *Pichia pastoris*: characterization and effect of its overexpression on the production of secreted, surface displayed and membrane proteins. *Microb Cell Fact.* 2010;9:49-60. doi 10.1186/1475-2859-9-49
- Han M., Wang W., Gong X., Zhou J., Xu C., Li Y. Increased expression of recombinant chitosanase by co-expression of Hac1p in the yeast *Pichia pastoris*. *Protein Pept Lett.* 2021;28(12):1434-1441. doi 10.2174/0929866528666211105111155
- Hartl F.U., Bracher A., Hayer-Hartl M. Molecular chaperones in protein folding and proteostasis. *Nature.* 2011;475(7356):324-332. doi 10.1038/nature10317
- Hatahet F., Ruddock L.W. Protein disulfide isomerase: a critical evaluation of its function in disulfide bond formation. *Antioxid Redox Signal.* 2009;11(11):2807-2850. doi 10.1089/ars.2009.2466
- Hendershot L.M., Buck T.M., Brodsky J.L. The essential functions of molecular chaperones and folding enzymes in maintaining endoplasmic reticulum homeostasis. *J Mol Biol.* 2024;436(14):168418. doi 10.1016/j.jmb.2023.168418
- Hernández-Elvira M., Torres-Quiroz F., Escamilla-Ayala A., Domínguez-Martin E., Escalante R., Kawasaki L., Ongay-Larios L., Coria R. The unfolded protein response pathway in the yeast *Kluyveromyces fragilis*. A comparative view among yeast species. *Cells.* 2018;7(8):106. doi 10.3390/cells7080106
- Herscovics A. Processing glycosidases of *Saccharomyces cerevisiae*. *Biochim Biophys Acta.* 1999;1426(2):275-285. doi 10.1016/S0304-4165(98)00129-9
- Huang J., Zhao Q., Chen L., Zhang C., Bu W., Zhang X., Zhang K., Yang Z. Improved production of recombinant *Rhizomucor miehei* lipase by coexpressing protein folding chaperones in *Pichia pastoris*, which triggered ER stress. *Bioengineered.* 2020;11(1):375-385. doi 10.1080/21655979.2020.1738127
- Huang M., Gao Y., Zhou X., Zhang Y., Cai M. Regulating unfolded protein response activator HAC1p for production of thermostable raw-starch hydrolyzing α -amylase in *Pichia pastoris*. *Bioprocess Biosyst Eng.* 2017;40(3):341-350. doi 10.1007/s00449-016-1701-y
- Inan M., Fanders S.A., Zhang W., Hotez P.J., Zhan B., Meagher M.M. Saturation of the secretory pathway by overexpression of a hookworm (*Necator americanus*) protein (Na-ASP1). In: Cregg J.M. (Ed.) *Pichia Protocols. Methods in Molecular Biology.* Vol. 389. Humana Press, 2007:65-76. doi 10.1007/978-1-59745-456-8_5
- Ishiwata-Kimata Y., Kimata Y. Fundamental and applicative aspects of the unfolded protein response in yeasts. *J Fungi (Basel).* 2023;9(10):989. doi 10.3390/jof9100989

- Ito Y., Ishigami M., Hashiba N., Nakamura Y., Terai G., Hasunuma T., Ishii J., Kondo A. Avoiding entry into intracellular protein degradation pathways by signal mutations increases protein secretion in *Pichia pastoris*. *Microb Biotechnol.* 2022;15(9):2364-2378. doi 10.1111/1751-7915.14061
- Kelleher D.J., Gilmore R. An evolving view of the eukaryotic oligosaccharyltransferase. *Glycobiology.* 2006;16(4):47R-62R. doi 10.1093/glycob/cwj066
- Khlebodarova T.M., Bogacheva N.V., Zadorozhny A.V., Bryanskaya A.V., Vasilieva A.R., Chesnokov D.O., Pavlova E.I., Peltek S.E. *Komagataella phaffii* as a platform for heterologous expression of enzymes used for industry. *Microorganisms.* 2024;12(2):346. doi 10.3390/microorganisms12020346
- Kim M.D., Han K.C., Kang H.A., Rhee S.K., Seo J.H. Coexpression of BIP increased antithrombotic hirudin production in recombinant *Saccharomyces cerevisiae*. *J Biotechnol.* 2003;101(1):81-87. doi 10.1016/s0168-1656(02)00288-2
- Kimmig P., Diaz M., Zheng J., Williams C.C., Lang A., Aragón T., Li H., Walter P. The unfolded protein response in fission yeast modulates stability of select mRNAs to maintain protein homeostasis. *eLife.* 2012;1:e00048. doi 10.7554/eLife.00048
- Korennykh A., Walter P. Structural basis of the unfolded protein response. *Annu Rev Cell Dev Biol.* 2012;28:251-277. doi 10.1146/annurev-cellbio-101011-155826
- Krshnan L., van de Weijer M.L., Carvalho P. Endoplasmic reticulum-associated protein degradation. *Cold Spring Harb Perspect Biol.* 2022;14(12):a041247. doi 10.1101/cshperspect.a041247
- Lee T.H., Bae Y.H., Kim M.D., Seo J.H. Overexpression of Hac1 gene increased levels of both intracellular and secreted human kringle fragment in *Saccharomyces cerevisiae*. *Process Biochem.* 2012;47(12):2300-2305. doi 10.1016/j.procbio.2012.09.006
- Lehle L., Strahl S., Tanner W. Protein glycosylation, conserved from yeast to man: a model organism helps elucidate congenital human diseases. *Angew Chem Int Ed Engl.* 2006;45(41):6802-6818. doi 10.1002/anie.200601645
- Li C., Lin Y., Zheng X., Pang N., Liao X., Liu X., Huang Y., Liang S. Combined strategies for improving expression of *Citrobacter amalonaticus* phytase in *Pichia pastoris*. *BMC Biotechnol.* 2015;15:88. doi 10.1186/s12896-015-0204-2
- Li F., Chen Y., Qi Q., Wang Y., Yuan L., Huang M., Elsemman I.E., Feizi A., Kerkhoven E.J., Nielsen J. Improving recombinant protein production by yeast through genome-scale modeling using proteome constraints. *Nat Commun.* 2022;13(1):2969. doi 10.1038/s41467-022-30689-7
- Lin Y., Feng Y., Zheng L., Zhao M., Huang M. Improved protein production in yeast using cell engineering with genes related to a key factor in the unfolded protein response. *Metab Eng.* 2023;77:152-161. doi 10.1016/j.ymben.2023.04.004
- Madhavan A., Arun K.B., Sindhu R., Krishnamoorthy J., Reshmy R., Sirohi R., Pugazhendhi A., Awasthi M.K., Szakacs G., Binod P. Customized yeast cell factories for biopharmaceuticals: from cell engineering to process scale up. *Microb Cell Fact.* 2021;20(1):124. doi 10.1186/s12934-021-01617-z
- Mizunaga T., Katakura Y., Miura T., Maruyama Y. Purification and characterization of yeast protein disulfide isomerase. *J Biochem.* 1990;108(5):846-851. doi 10.1093/oxfordjournals.jbchem.a123291
- Mori A., Hara S., Sugahara T., Kojima T., Iwasaki Y., Kawarasaki Y., Sahara T., Ohgiya S., Nakano H. Signal peptide optimization tool for the secretion of recombinant protein from *Saccharomyces cerevisiae*. *J Biosci Bioeng.* 2015;120(5):518-525. doi 10.1016/j.jbiosc.2015.03.003
- Mori K. Evolutionary aspects of the unfolded protein response. *Cold Spring Harb Perspect Biol.* 2022;14(12):a041262. doi 10.1101/cshperspect.a041262
- Niu Y., Zhang L., Yu J., Wang C.C., Wang L. Novel roles of the non-catalytic elements of yeast protein-disulfide isomerase in its interplay with endoplasmic reticulum oxidoreductin 1. *J Biol Chem.* 2016;291(15):8283-8294. doi 10.1074/jbc.M115.694257
- Núñez A., Dulude D., Jbel M., Rokeach L.A. Calnexin is essential for survival under nitrogen starvation and stationary phase in *Schizosaccharomyces pombe*. *PLoS One.* 2015;10(3):e0121059. doi 10.1371/journal.pone.0121059
- O'Keefe S., Pool M.R., High S. Membrane protein biogenesis at the ER: the highways and byways. *FEBS J.* 2022;289(22):6835-6862. doi 10.1111/febs.15905
- Omkar S., Mitchem M.M., Hoskins J.R., Shrader C., Kline J.T., Nitika, Fornelli L., Wickner S., Truman A.W. Acetylation of the yeast Hsp40 chaperone protein Ydj1 fine-tunes proteostasis and translational fidelity. *PLoS Genet.* 2024;20(12):e1011338. doi 10.1371/journal.pgen.1011338
- Palma A., Rettenbacher L.A., Moilanen A., Saaranen M., Pacheco-Martinez C., Gasser B., Ruddock L. Biochemical analysis of *Komagataella phaffii* oxidative folding proposes novel regulatory mechanisms of disulfide bond formation in yeast. *Sci Rep.* 2023;13(1):14298. doi 10.1038/s41598-023-41375-z
- Parlati F., Dominguez M., Bergeron J.J., Thomas D.Y. *Saccharomyces cerevisiae* CNE1 encodes an endoplasmic reticulum (ER) membrane protein with sequence similarity to calnexin and calreticulin and functions as a constituent of the ER quality control apparatus. *J Biol Chem.* 1995;270(1):244-253. doi 10.1074/jbc.270.1.244
- Parodi A.J. Protein glucosylation and its role in protein folding. *Annu Rev Biochem.* 2000;69:69-93. doi 10.1146/annurev.biochem.69.1.69
- Payne T., Finnis C., Evans L.R., Mead D.J., Avery S.V., Archer D.B., Sleep D. Modulation of chaperone gene expression in mutagenized *Saccharomyces cerevisiae* strains developed for recombinant human albumin production results in increased production of multiple heterologous proteins. *Appl Environ Microbiol.* 2008;74(24):7759-7766. doi 10.1128/AEM.01178-08
- Pfeffer M., Maurer M., Stadlmann J., Grass J., Delic M., Altmann F., Mattanovich D. Intracellular interactome of secreted antibody Fab fragment in *Pichia pastoris* reveals its routes of secretion and degradation. *Appl Microbiol Biotechnol.* 2012;93(6):2503-2512. doi 10.1007/s00253-012-3933-3
- Pickart C.M. Mechanisms underlying ubiquitination. *Annu Rev Biochem.* 2001;70:503-533. doi 10.1146/annurev.biochem.70.1.503
- Preston G.M., Brodsky J.L. The evolving role of ubiquitin modification in endoplasmic reticulum-associated degradation. *Biochem J.* 2017;474(4):445-469. doi 10.1042/BCJ20160582
- Qi Q., Li F., Yu R., Engqvist M.K.M., Siewers V., Fuchs J., Nielsen J. Different routes of protein folding contribute to improved protein production in *Saccharomyces cerevisiae*. *mBio.* 2020;11(6):e02743-20. doi 10.1128/mBio.02743-20
- Raschmanová H., Weninger A., Knejzlík Z., Melzoch K., Kovar K. Engineering of the unfolded protein response pathway in *Pichia pastoris*: enhancing production of secreted recombinant proteins. *Appl Microbiol Biotechnol.* 2021;105(11):4397-4414. doi 10.1007/s00253-021-11336-5
- Robinson A.S., Hines V., Wittrup K.D. Protein disulfide isomerase overexpression increases secretion of foreign proteins in *Saccharomyces cerevisiae*. *Nat Biotechnol.* 1994;12(4):381-384. doi 10.1038/nbt0494-381
- Ruger-Herreros C., Svoboda L., Mogk A., Bukau B. Role of J-domain proteins in yeast physiology and protein quality control. *J Mol Biol.* 2024;436(14):168484. doi 10.1016/j.jmb.2024.168484
- Ruggiano A., Foresti O., Carvalho P. Quality control: ER-associated degradation: protein quality control and beyond. *J Cell Biol.* 2014;204(6):869-879. doi 10.1083/jcb.201312042
- Saibil H. Chaperone machines for protein folding, unfolding and disaggregation. *Nat Rev Mol Cell Biol.* 2013;14(10):630-642. doi 10.1038/nrm3658
- Sallada N.D., Harkins L.E., Berger B.W. Effect of gene copy number and chaperone coexpression on recombinant hydrophobin HFBI biosurfactant production in *Pichia pastoris*. *Biotechnol Bioeng.* 2019;116(8):2029-2040. doi 10.1002/bit.26982
- Schlenstedt G., Harris S., Risse B., Lill R., Silver P.A. A yeast DnaJ homologue, Scj1p, can function in the endoplasmic reticulum with

- BiP/Kar2p via a conserved domain that specifies interactions with Hsp70s. *J Cell Biol.* 1995;129(4):979-988. doi 10.1083/jcb.129.4.979
- Schroder M., Clark R., Kaufman R.J. *IRE1*- and *HAC1*-independent transcriptional regulation in the unfolded protein response of yeast. *Mol Microbiol.* 2003;49(3):591-606. doi 10.1046/j.1365-2958.2003.03585.x
- Schuck S., Prinz W.A., Thorn K.S., Voss C., Walter P. Membrane expansion alleviates endoplasmic reticulum stress independently of the unfolded protein response. *J Cell Biol.* 2009;187(4):525-536. doi 10.1083/jcb.200907074
- Schwarz D.S., Blower M.D. The endoplasmic reticulum: structure, function and response to cellular signaling. *Cell Mol Life Sci.* 2016; 73(1):79-94. doi 10.1007/s00018-015-2052-6
- Sevier C.S., Kaiser C.A. Disulfide transfer between two conserved cysteine pairs imparts selectivity to protein oxidation by Ero1. *Mol Biol Cell.* 2006;17(5):2256-2266. doi 10.1091/mbc.e05-05-0417
- Sheng J., Flick H., Feng X. Systematic optimization of protein secretory pathways in *Saccharomyces cerevisiae* to increase expression of hepatitis B small antigen. *Front Microbiol.* 2017;8:875. doi 10.3389/fmicb.2017.00875
- Shusta E.V., Raines R.T., Plückthun A., Wittup K.D. Increasing the secretory capacity of *Saccharomyces cerevisiae* for production of single-chain antibody fragments. *Nat Biotechnol.* 1998;16(8):773-777. doi 10.1038/nbt0898-773
- Singhvi P., Saneja A., Ahuja R., Panda A.K. Solubilization and refolding of variety of inclusion body proteins using a novel formulation. *Int J Biol Macromol.* 2021;193(Pt.B):2352-2364. doi 10.1016/j.ijbiomac.2021.11.068
- Smith J.D., Tang B.C., Robinson A.S. Protein disulfide isomerase, but not binding protein, overexpression enhances secretion of a non-disulfide-bonded protein in yeast. *Biotechnol Bioeng.* 2004;85(3): 340-350. doi 10.1002/bit.10853
- Steel G.J., Fullerton D.M., Tyson J.R., Stirling C.J. Coordinated activation of Hsp70 chaperones. *Science.* 2004;303(5654):98-101. doi 10.1126/science.1092287
- Thak E.J., Yoo S.J., Moon H.Y., Kang H.A. Yeast synthetic biology for designed cell factories producing secretory recombinant proteins. *FEMS Yeast Res.* 2020;20(2):foaa009. doi 10.1093/femsyr/foaa009
- Thibault G., Ng D.T.W. The endoplasmic reticulum-associated degradation pathways of budding yeast. *Cold Spring Harb Perspect Biol.* 2012;4(12):a013193. doi 10.1101/cshperspect.a013193
- Travers K.J., Patil C.K., Wodicka L., Lockhart D.J., Weissman J.S., Walter P. Functional and genomic analyses reveal an essential coordination between the unfolded protein response and ER-associated degradation. *Cell.* 2000;101(3):249-258. doi 10.1016/s0092-8674(00)80835-1
- Tsuda M., Nonaka K. Recent progress on heterologous protein production in methylotrophic yeast systems. *World J Microbiol Biotechnol.* 2024;40(7):200. doi 10.1007/s11274-024-04008-9
- Valkonen M., Penttilä M., Saloheimo M. Effects of inactivation and constitutive expression of the unfolded-protein response pathway on protein production in the yeast *Saccharomyces cerevisiae*. *Appl Environ Microbiol.* 2003;69(4):2065-2072. doi 10.1128/aem.69.4.2065-2072.2003
- Wang L., Wang C.C. Oxidative protein folding fidelity and redox status in the endoplasmic reticulum. *Trends Biochem Sci.* 2023;48(1):40-52. doi 10.1016/j.tibs.2022.06.011
- Ware F.E., Vassilakos A., Peterson P.A., Jackson M.R., Lehman M.A., Williams D.B. The molecular chaperone calnexin binds Glc1Man9GlcNAc2 oligosaccharide as an initial step in recognizing unfolded glycoproteins. *J Biol Chem.* 1995;270(9):4697-4704. doi 10.1074/jbc.270.9.4697
- Xia X. Translation control of *HAC1* by regulation of splicing in *Saccharomyces cerevisiae*. *Int J Mol Sci.* 2019;20(12):2860. doi 10.3390/ijms20122860
- Xu C., Ng D.T.W. Glycosylation-directed quality control of protein folding. *Nat Rev Mol Cell Biol.* 2015;16:742-752. doi 10.1038/nrm4073
- Xu X., Kanbara K., Azakam H., Kato A. Expression and characterization of *Saccharomyces cerevisiae* Cne1p, a calnexin homologue. *J Biochem.* 2004;135(5):615-618. doi 10.1093/jb/mvh074
- Yamaguchi H., Miyazaki M. Refolding techniques for recovering biologically active recombinant proteins from inclusion bodies. *Bio-molecules.* 2014;4(1):235-251. doi 10.3390/biom4010235
- Yang J., Lu Z., Chen J., Chu P., Cheng Q., Liu J., Ming F., Huang C., Xiao A., Cai H., Zhang L. Effect of cooperation of chaperones and gene dosage on the expression of porcine PGLYRP-1 in *Pichia pastoris*. *Appl Microbiol Biotechnol.* 2016;100(12):5453-5465. doi 10.1007/s00253-016-7372-4
- Zahrl R.J., Gasser B., Mattanovich D., Ferrer P. Detection and elimination of cellular bottlenecks in protein-producing yeasts. In: Gasser B., Mattanovich D. (Eds) Recombinant Protein Production in Yeast. Methods in Molecular Biology. Vol. 1923. Humana Press, 2019;75-95. doi 10.1007/978-1-4939-9024-5_2
- Zahrl R.J., Prielhofer R., Ata Ö., Baumann K., Mattanovich D., Gasser B. Pushing and pulling proteins into the yeast secretory pathway enhances recombinant protein secretion. *Metab Eng.* 2022;74: 36-48. doi 10.1016/j.ymben.2022.08.010
- Zahrl R.J., Prielhofer R., Burgard J., Mattanovich D., Gasser B. Synthetic activation of yeast stress response improves secretion of recombinant proteins. *N Biotechnol.* 2023;73:19-28. doi 10.1016/j.nbt.2023.01.001
- Zha J., Liu D., Ren J., Liu Z., Wu X. Advances in metabolic engineering of *Pichia pastoris* strains as powerful cell factories. *J Fungi (Basel).* 2023;9(10):1027. doi 10.3390/jof9101027

Conflict of interest. The authors declare no conflict of interest.

Received September 2, 2025. Revised November 25, 2025. Accepted November 25, 2025.