

## SUPPLEMENTARY MATERIALS

to the article: V.Y. Vavilova, I. Konopatskaia, S.L. Luzyanin, M. Woyciechowski, A.G. Blinov  
 "Parasites of the genus *Nosema*, *Critidilia* and *Lotmaria* in the honeybee and bumblebee populations:  
 a case study in India"

**Table S1.** The literature references about microsporidian parasites, *N. apis* and *N. ceranae*, in honeybees

Country	<i>N. apis</i> infection	<i>N. ceranae</i> infection	References
Argentina	NI	+	Branchiccela et al., 2017
Australia	+	+	Rice, 2001; Dosselli et al., 2016; Sagastume et al., 2016
Belgium	+	+	Ravoet et al., 2013
Brazil	+	+	Gatehouse, Malone, 1999; Klee et al., 2007; Teixeira et al., 2013; Branchiccela et al., 2017
Canada	+	+	Gatehouse, Malone, 1999; Rice, 2001; Hamiduzzaman et al., 2010; Shutler et al., 2014; Emsen et al., 2016; Branchiccela et al., 2017
China	+	+	Chen et al., 2009a; Yang et al., 2013
Croatia		+	Branchiccela et al., 2017
Czech Republic	+	+	Hubert et al., 2017
Denmark	+	+	Klee et al., 2007
Finland	+	+	Gatehouse, Malone, 1999; Klee et al., 2007
France	NI	+	Sagastume et al., 2016; Branchiccela et al., 2017
Germany	+	+	Klee et al., 2007; Sagastume et al., 2016; Wells et al., 2016; Gisder et al., 2017
Greece	-	+	Klee et al., 2007; Bacandritsos et al., 2010
Hungary	+	+	Klee et al., 2007; Csáki et al., 2015
Indonesia	+	-	Rice, 2001
Iran	-	+	Nabian et al., 2011
Ireland	+	+	Klee et al., 2007
Israel	+	-	Gatehouse, Malone, 1999
Italy	+	+	Gatehouse, Malone, 1999; Klee et al., 2007; Porrini et al., 2016; Branchiccela et al., 2017
Japan	+	+	Chen et al., 2009a; Morimoto et al., 2013
Kenya	+	-	Muli et al., 2014
Mexico	NI	+	Guerrero-Molina et al., 2016
Mongolia	-	+	Tsevegmid et al., 2016
New Zealand	+	+	Gatehouse, Malone, 1998, 1999; Rice, 2001; Klee et al., 2007
Poland	+	+	Ptaszyńska et al., 2016
Portugal		+	Branchiccela et al., 2017
Russia	+	+	Zinatullina et al., 2013; Ostroverkhova et al., 2016a,b
Saudi Arabia	-	+	Ansari et al., 2017
Serbia	-	+	Klee et al., 2007; Stevanovic et al., 2016
Slovakia	NI	+	Sagastume et al., 2016
Slovenia	NI	+	Sagastume et al., 2016; Branchiccela et al., 2017
Spain	+	+	Klee et al., 2007; Higes et al., 2010; Jara et al., 2012, 2015; Botías et al., 2013; Cepero et al., 2014; Sagastume et al., 2016; Branchiccela et al., 2017
Sweden	+	+	Gatehouse, Malone, 1999; Klee et al., 2007; Forsgren, Fries, 2013
Switzerland	+	+	Gatehouse, Malone, 1999; Dainat et al., 2012; Sagastume et al., 2016; Branchiccela et al., 2017; Tritschler et al., 2017
Taiwan	+	+	Chen et al., 2009a
the Netherlands	NI	+	Branchiccela et al., 2017
Turkey	+	+	Whitaker et al., 2011; İvgin Tunca et al., 2016; Branchiccela et al., 2017
UK	+	+	Gatehouse, Malone, 1999; Klee et al., 2007; Budge et al., 2015; Roberts et al., 2015
Uruguay		+	Branchiccela et al., 2017
USA	+	+	Gatehouse, Malone, 1998; Klee et al., 2007; Chen et al., 2008, 2009b; Traver, Fell, 2011; Traver et al., 2012; Zheng et al., 2015; Cavigli et al., 2016
Vietnam	-	+	Klee et al., 2007
Zimbabwe	+	-	Fries et al., 2003

**Table S2.** The literature references about trypanosomatid parasites, *C. mellifcae* and *L. passim*, in honeybees

Country	<i>C. mellifcae</i> infection	<i>L. passim</i> infection	References
Australia	+	NI	Langridge, McGhee, 1967
Belgium	+	+	Ravoet et al., 2015; Schwarz et al., 2015
Japan	-	+	Ravoet et al., 2015
Serbia	-	+	Stevanovic et al., 2016
Spain	+	NI	Cepero et al., 2014
Switzerland	-	+	Ravoet et al., 2015; Tritschler et al., 2017
USA	+	NI	Runckel et al., 2011; Cornman et al., 2012

**Table S3.** The literature references about trypanosomatid parasites,  
*C. bombi*, *C. expoeki* and *C. mexicana*, in bumblebees

Country	<i>C. bombi</i> infection	<i>C. expoeki</i> infection	<i>C. mexicana</i> infection	References
Argentina	+	NI	NI	Schmid-Hempel et al., 2014
Argentina	+	NI	NI	Plischuk et al., 2017
Belgium	+	NI	NI	Meeus et al., 2010
Chile	+	NI	NI	Schmid-Hempel et al., 2014
England	+	NI	NI	Graystock et al., 2014
Germany	+	NI	NI	Popp et al., 2012
Germany	+	NI	NI	Erler et al., 2012
Italy	+	NI	NI	Lipa, Triggiani, 1988
Mexico	-	+	+	Gallot-Lavallée et al., 2016
Russia	+	NI	NI	Gorbunov, 1987
Switzerland	+	+	NI	Schmid-Hempel, Reber Funk, 2004; Schmid-Hempel, Tognazzo, 2010; Salathé, Schmid-Hempel, 2011

## References

- Ansari M.J., Al-Ghamdi A., Nuru A., Khan K.A., Alattal J. Geographical distribution and molecular detection of *Nosema ceranae* from indigenous honey bees of Saudi Arabia. *Saudi J. Biol. Sci.* 2017;24(5):983-991.
- Bacandritsos N., Granato A., Budge G., Papanastasiou I., Rojiani E., Caldon M., Falcaro C., Gallina A., Mutinelli F. Sudden deaths and colony population decline in Greek honey bee colonies. *J. Invertebr. Pathol.* 2010;105(3):335-340.
- Botías C., Martín-Hernández R., Barrios L., Meana S., Higes M. *Nosema* spp. infection and its negative effects on honey bees (*Apis mellifera iberiensis*) at the colony level. *Vet. Res.* 2013;44(1):25.
- Branchiccela B., Arredondo D., Higes M., Invernizzi C., Martín-Hernández R., Tomasco I., Zunino P., Antúnez K. Characterization of *Nosema ceranae* genetic variants from different geographic origins. *Microb. Ecol.* 2017;73(4):978-987.
- Budge G.E., Pietravalle S., Brown M., Laurenson L., Jones B., Tomkies V., Delaplane K.S. Pathogens as predictors of honey bee colony strength in England and Wales. *PLoS ONE.* 2015;10(7):e0133228.
- Cavigli I., Daughenbaugh K.F., Martin M., Lerch M., Banner K., Garcia E., Brutscher L.M., Flenniken M.L. Pathogen prevalence and abundance in honey bee colonies involved in almond pollination. *Apidologie.* 2016;47:251-266.
- Cepero A., Ravoet J., Gómez-Moracho T., Bernal J.L., Del Nozal M.J., Bartolomé C., Maside X., Meana A., González-Porto A.V., de Graaf D.C., Martín-Hernández R., Higes M. Holistic screening of collapsing honey bee colonies in Spain: a case study. *BMC Res. Notes.* 2014;7:649.
- Chen Y., Evans J.D., Smith I.B., Pettis J.S. *Nosema ceranae* is a long-present and wide-spread microsporidian infection of the European honey bee (*Apis mellifera*) in the United States. *J. Invertebr. Pathol.* 2008;97(2):186-188.
- Chen Y., Evans J.D., Zhou L., Boncristiani H., Kimura K., Xiao T., Litkowski A.M., Pettis J.S. Asymmetrical coexistence of *Nosema ceranae* and *Nosema apis* in honey bees. *J. Invertebr. Pathol.* 2009a;101(3):204-9.
- Chen Y.P., Evans J.D., Murphy C., Gutell R., Zuker M., Gundensen-Rindal D., Pettis J.S. Morphological, molecular, and phylogenetic characterization of *Nosema ceranae*, a microsporidian parasite isolated from the European honey bee, *Apis mellifera*. *J. Eukaryot. Microbiol.* 2009b;56(2):142-147.
- Cornman R.S., Tarpy D.R., Chen Y., Jeffreys L., Lopez D., Pettis J.S., VanEngelsdorp D., Evans J.D. Pathogen webs in collapsing honey bee colonies. *PLoS ONE.* 2012;7:e43562.
- Csáki T., Heltai M., Markolt F., Kovács B., Békési L., Ladányi M., Péntek-Zakar E., Meana A., Botías C., Martín-Hernández R., Higes M. Permanent prevalence of *Nosema ceranae* in honey bees (*Apis mellifera*) in Hungary. *Acta Vet. Hung.* 2015;63(3):358-369.
- Dainat B., Evans J.D., Chen Y.P., Gauthier L., Neumann P. Predictive markers of honey bee colony collapse. *PLoS ONE.* 2012;7(2):e32151.
- Dosselli R., Grassl J., Carson A., Simmons L.W., Baer B. Flight behaviour of honey bee (*Apis mellifera*) workers is altered by initial infections of the fungal parasite *Nosema apis*. *Sci. Rep.* 2016;6:36649.
- Emsen B., Guzman-Novoa E., Hamiduzzaman M.M., Eccles L., Lacey B., Ruiz-Pérez R.A., Nasr M. Higher prevalence and levels of *Nosema ceranae* than *Nosema apis* infections in Canadian honey bee colonies. *Parasitol. Res.* 2016;115(1):175-181.
- Erler S., Popp M., Wolf S., Latorff H.M.G. Sex, horizontal transmission, and multiple hosts prevent local adaptation of *Crithidia bombi*, a parasite of bumblebees (*Bombus* spp.). *Ecol. Evol.* 2012;2(5):930-940.
- Forsgren E., Fries I. Temporal study of *Nosema* spp. in a cold climate. *Environ. Microbiol. Rep.* 2013;5(1):78-82.
- Fries I., Slemenda S.B., da Silva A., Pieniazek N.J. African honey bees (*Apis mellifera scutellata*) and nosema (*Nosema apis*) infections. *J. Apicult. Res.* 2003;42:13-15.
- Gallot-Lavallée M., Schmid-Hempel R., Vandame R., Vergara C.H., Schmid-Hempel P. Large scale patterns of abundance and distribution of parasites in Mexican bumblebees. *J. Invertebr. Pathol.* 2016;133:73-82.
- Gatehouse H.S., Malone L.A. Genetic variability among *Nosema apis* isolates. *J. Apicult. Res.* 1999;38:79-85.
- Gatehouse H.S., Malone L.A. The ribosomal RNA gene region of *Nosema apis* (microspora): DNA sequence for small and large sub-unit rRNA genes and evidence of a large tandem repeat unit size. *J. Invertebr. Pathol.* 1998;71:97-105.
- Gisder S., Schüller V., Horchler L.L., Groth D., Genersch E. Long-term temporal trends of *Nosema* spp. infection prevalence in north-east Germany: continuous spread of *Nosema ceranae*, an emerging pathogen of honey bees (*Apis mellifera*), but no general replacement of *Nosema apis*. *Front. Cell. Infect. Microbiol.* 2017;7:301.
- Gorbunov P.S. Endoparasitic flagellates of the genus *Crithidia* (Trypanosomatidae, Zoomastigophorea) from the digestive tract of bumblebees. *Russ. J. Zool.* 1987;66(12):1775-1780. (in Russian)
- Graystock P., Goulson D., Hughes W.O. The relationship between managed bees and the prevalence of parasites in bumblebees. *PeerJ.* 2014;2:e522.
- Guerrero-Molina C., Correa-Benítez A., Hamiduzzaman M.M., Guzman-Novoa E. *Nosema ceranae* is an old resident of honey bee (*Apis mellifera*) colonies in Mexico, causing infection levels of one million spores per bee or higher during summer and fall. *J. Invertebr. Pathol.* 2016;141:38-40.
- Hamiduzzaman M.M., Guzman-Novoa E., Goodwin P.H. A multiplex PCR assay to diagnose and quantify *Nosema* infections in honey bees (*Apis mellifera*). *J. Invertebr. Pathol.* 2010;105(2):151-155.
- Higes M., Martín-Hernández R., Martínez-Salvador A., Garrido-Bailón E., González-Porto A.V., Meana A., Bernal J.L., Del Nozal M.J., Bernal J. A preliminary study of the epidemiological factors related to honey bee colony loss in Spain. *Environ. Microbiol. Rep.* 2010;2(2):243-250.
- Hubert J., Bicianova M., Ledvinka O., Kamler M., Lester P.J., Nesvorna M., Kopecky J., Erban T. Changes in the bacteriome of honey bees associated with the parasite *Varroa destructor*, and pathogens *Nosema* and *Lotmaria passim*. *Microb. Ecol.* 2017;73(3):685-698.
- Ivgin Tunca R., Oskay D., Gosterit A., Tekin O.K. Does *Nosema ceranae* wipe out *Nosema apis* in Turkey? *Iran J. Parasitol.* 2016;11(2):259-264.

- Jara L., Cepero A., Garrido-Bailón E., Martín-Hernández R., Higes M., De la Rúa P. Linking evolutionary lineage with parasite and pathogen prevalence in the Iberian honey bee. *J. Invertebr. Pathol.* 2012;110(1):8-13.
- Jara L., Muñoz I., Cepero A., Martín-Hernández R., Serrano J., Higes M., De la Rúa P. Stable genetic diversity despite parasite and pathogen spread in honey bee colonies. *Naturwissenschaften*. 2015;102(9-10):53.
- Klee J., Besana A.M., Genersch E., Gisder S., Nanetti A., Tam D.Q., Chinh T.X., Puerta F., Ruz J.M., Kryger P., Message D., Hatjina F., Korpela S., Fries I., Paxton R.J. Widespread dispersal of the microsporidian *Nosema ceranae*, an emergent pathogen of the western honey bee, *Apis mellifera*. *J. Invertebr. Pathol.* 2007;96(1):1-10.
- Langridge D.F., McGhee R.B. *Crithidia mellifica* n. sp.: An acidophilic trypanosomatid of honey bee *Apis mellifera*. *J. Protozool.* 1967;14:485-487.
- Lipa J.J., Triggiani O. *Crithidia bombi* sp. n. a new flagellated parasite of a bumble bee *Bombus terrestris* L. (Hymenoptera, Apidae). *Acta Protozool.* 1988;27:287-290.
- Meeus I., De Graaf D.C., Jans K., Smagghe G. Multiplex PCR detection of slowly-evolving trypanosomatids and neogregarines in bumblebees using broad-range primers. *J. Appl. Microbiol.* 2010;109:107-115.
- Morimoto T., Kojima Y., Yoshiyama M., Kimura K., Yang B., Peng G., Kadokawa T. Molecular detection of protozoan parasites infecting *Apis mellifera* colonies in Japan. *Environ. Microbiol. Rep.* 2013;5(1):74-77.
- Muli E., Patch H., Frazier M., Frazier J., Torto B., Baumgarten T., Kilonzo J., Kimani J.N., Mumoki F., Masiga D., Tumlinson J., Grozinger K. Evaluation of the distribution and impacts of parasites, pathogens, and pesticides on honey bee (*Apis mellifera*) populations in East Africa. *PLoS ONE*. 2014;9(4):e94459.
- Nabian S., Ahmadi K., Nazem Shirazi M.H., Gerami Sadeghian A. First detection of *Nosema ceranae*, a microsporidian protozoa of European honeybees (*Apis mellifera*) in Iran. *Iran J. Parasitol.* 2011;6(3):89-95.
- Ostroverkhova N.V., Konusova O.L., Kucher A.N., Golubeva E.P., Kireeva T.N., Pogor Y.L. Nosemosis type C in the Tomsk region. *Beekeeping*. 2016a;8. (in Russian)
- Ostroverkhova N.V., Konusova O.L., Kucher A.N., Simakova A.V., Golubeva E.P., Kireeva T.N., Sharakhov I.V. Infestation of honey-bee (*Apis mellifera*) families by Microsporidia of *Nosema* genus (Microsporidia) at the apiaries of Tomsk region. *Parasitology*. 2016b;50:197-210. (in Russian)
- Plischuk S., Antúnez K., Harambou M., Minardi G.M., Lange C.E. Long-term prevalence of the protists *Crithidia bombi* and *Apicystis bombi* and detection of the microsporidium *Nosema bombi* in invasive bumble bees. *Environ. Microbiol. Rep.* 2017;9(2):169-173.
- Popp M., Erler S., Lattorff H.M.G. Seasonal variability of prevalence and occurrence of multiple infections shape the population structure of *Crithidia bombi*, an intestinal parasite of bumblebees (*Bombus* spp.). *Microbiologyopen*. 2012;1(4):362-372.
- Porrini C., Mutinelli F., Bortolotti L., Granato A., Laurenson L., Roberts K., Gallina A., Silvester N., Medrzycki P., Renzi T., Sgolastra F., Lodesani M. The status of honey bee health in Italy: results from the nationwide bee monitoring network. *PLoS ONE*. 2016;11(5):e0155411.
- Ptaszyńska A.A., Paleolog J., Borsuk G. *Nosema ceranae* infection promotes proliferation of yeasts in honey bee intestines. *PLoS ONE*. 2016;11(10):e0164477.
- Ravoet J., Maharramov J., Meeus I., De Smet L., Wenseleers T., Smagghe G., de Graaf D.C. Comprehensive bee pathogen screening in Belgium reveals *Crithidia mellifica* as a new contributory factor to winter mortality. *PLoS ONE*. 2013;8:e72443.
- Ravoet J., Schwarz R.S., Descamps T., Yañez O., Tozkar C.O., Martin-Hernandez R., Bartolomé C., De Smet L., Higes M., Wenseleers T., Schmid-Hempel R., Neumann P., Kadokawa T., Evans J.D., de Graaf D.C. Differential diagnosis of the honey bee trypanosomatids *Crithidia mellifica* and *Lotmaria passim*. *J. Invertebr. Pathol.* 2015;130:21-27.
- Rice R.N. *Nosema* Disease in Honeybees – Genetic Variation and Control. Australia: Rural Industries Research and Development Corporation, 2001. Publ. No. 01/46.
- Roberts K.E., Evison S.E., Baer B., Hughes W.O. The cost of promiscuity: sexual transmission of *Nosema* microsporidian parasites in polyandrous honey bees. *Sci. Rep.* 2015;5:10982.
- Runckel C., Flenniken M.L., Engel J.C., Ruby J.G., Ganem D., Andino R., DeRisi J.L. Temporal analysis of the honey bee microbiome reveals four novel viruses and seasonal prevalence of known viruses, *Nosema*, and *Crithidia*. *PLoS ONE*. 2011;6:e20656.
- Sagastume S., Martín-Hernández R., Higes M., Henriques-Gil N. Genotype diversity in the honey bee parasite *Nosema ceranae*: multi-strain isolates, cryptic sex or both? *BMC Evol. Biol.* 2016;16(1):216.
- Salathé R.M., Schmid-Hempel P. The genotypic structure of a multi-host bumblebee parasite suggests a role for ecological niche overlap. *PLoS ONE*. 2011;6(8):e22054.
- Schmid-Hempel P., Reber Funk C. The distribution of genotypes of the trypanosome parasite, *Crithidia bombi*, in populations of its host, *Bombus terrestris*. *Parasitology*. 2004;129:147-158.
- Schmid-Hempel R., Eckhardt M., Goulson D., Heinzmann D., Lange C., Plischuk S., Escudero L.R., Salathé R., Scriven J.J., Schmid-Hempel P. The invasion of southern South America by imported bumblebees and associated parasites. *J. Anim. Ecol.* 2014;83: 823-837.
- Schmid-Hempel R., Tognazzo M. Molecular divergence defines two distinct lineages of *Crithidia bombi* (Trypanosomatidae), parasites of bumblebees. *J. Eukaryot. Microbiol.* 2010;57:337-345.
- Schwarz R.S., Bauchan G.R., Murphy C.A., Ravoet J., De Graaf D.C., Evans J.D. Characterization of two species of trypanosomatidae from the honey bee *Apis mellifera*: *Crithidia mellifica* Langridge and McGhee, and *Lotmaria passim* n. gen., n. sp. *J. Eukaryot. Microbiol.* 2015;62:567-583.
- Shutler D., Head K., Burgher-MacLellan K.L., Colwell M.J., Levitt A.L., Ostiguy N., Williams G.R. Honey bee *Apis mellifera* parasites in the absence of *Nosema ceranae* fungi and *Varroa destructor* mites. *PLoS ONE*. 2014;9(6):e98599.
- Stevanovic J., Schwarz R.S., Vejnovic B., Evans J.D., Irwin R.E., Glavinic U., Stanimirovic Z. Species-specific diagnostics of *Apis mellifera* trypanosomatids: A nine-year survey (2007–2015) for trypanosomatids and microsporidians in Serbian honey bees. *J. Invertebr. Pathol.* 2016;139:6-11.

- Teixeira E.W., Santos L.G., Sattler A., Message D., Alves M.L., Martins M.F., Grassi-Sella M.L., Franco T.M. *Nosema ceranae* has been present in Brazil for more than three decades infecting Africanized honey bees. *J. Invertebr. Pathol.* 2013;114(3):250-254.
- Traver B.E., Fell R.D. Prevalence and infection intensity of *Nosema* in honey bee (*Apis mellifera* L.) colonies in Virginia. *J. Invertebr. Pathol.* 2011;107(1):43-49.
- Traver B.E., Williams M.R., Fell R.D. Comparison of within hive sampling and seasonal activity of *Nosema ceranae* in honey bee colonies. *J. Invertebr. Pathol.* 2012;109(2):187-193.
- Tritschler M., Retschnig G., Yañez O., Williams G.R., Neumann P. Host sharing by the honey bee parasites *Lotmaria passim* and *Nosema ceranae*. *Ecol. Evol.* 2017;7(6):1850-1857.
- Tsevegmid K., Neumann P., Yañez O. The honey bee pathosphere of Mongolia: European viruses in Central Asia. *PLoS ONE*. 2016; 11(3):e0151164.
- Wells T., Wolf S., Nicholls E., Groll H., Lim K.S., Clark S.J., Swain J., Osborne J.L., Haughton A.J. Flight performance of actively foraging honey bees is reduced by a common pathogen. *Environ. Microbiol. Rep.* 2016;8(5):728-737.
- Whitaker J., Szalanski A.L., Kence M. Molecular detection of *Nosema ceranae* and *N. apis* from Turkish honey bees. *Apidologie*. 2011;42(2):174-180.
- Yang B., Peng G., Li T., Kadokawa T. Molecular and phylogenetic characterization of honey bee viruses, *Nosema* microsporidia, protozoan parasites, and parasitic mites in China. *Ecol. Evol.* 2013;3(2):298-311.
- Zheng H.Q., Gong H.R., Huang S.K., Sohr A., Hu F.L., Chen Y.P. Evidence of the synergistic interaction of honey bee pathogens *Nosema ceranae* and *Deformed wing virus*. *Vet. Microbiol.* 2015;177(1-2):1-6.
- Zinatullina Z.Y., Zhigileva O.N., Ignatieva A.N., Tokarev Y.S. Differential diagnostics of honeybee nosemosis pathogen. *Sci. Notes Kazan SAVM*. 2013;2:190-194. (in Russian)